

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:

(11) International Publication Number:

WO 99/51639

C07K 14/705, A61K 38/17, C07K 16/28

(43) International Publication Date:

14 October 1999 (14.10.99)

(21) International Application Number:

PCT/SE99/00544

(22) International Filing Date:

31 March 1999 (31.03.99)

(30) Priority Data:

9801164-6 9900319-6 2 April 1998 (02.04.98)

28 January 1999 (28.01.99)

SE SE

(71) Applicant (for all designated States except US): ACTIVE BIOTECH AB [SE/SE]; Scheelevägen 22, S-220 07 Lund

(72) Inventor; and

(75) Inventor/Applicant US LUND-(for only): GREN-ÅKERLUND, Evy [SE/SE]; Trollsjövägen 165, S-237 33 Bjärred (SE).

(74) Agent: AWAPATENT AB; P.O. Box 5117, S-200 71 Malmö

(81) Designated States: AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).


#### Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

## (54) Title: AN INTEGRIN HETERODIMER AND A SUBUNIT THEREOF

A recombinant or isolated integrin heterodimer comprising a novel subunit a 10 in association with a subunit  $\beta$  is described. The  $\alpha$  10 integrin may be purified from bovine chondrocytes on a collagen-type-II affinity column. The integrin or the subunit  $\alpha 10$  can be used as marker or target of all types of cells, e.g. of chondrocytes, osteoblasts and fibroblasts. The integrin or subunit a 10 thereof can be used as marker or target in different physiological or therapeutic methods. They can also be used as active ingredients in pharmaceutical compositions and vaccines.



BEST AVAILABLE COPY

# FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK .	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Моласо	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	<b>IS</b>	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam-
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland	•	
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Pederation		
DE	Germany	น	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## AN INTEGRIN HETERODIMER AND A SUBUNIT THEREOF

#### FIELD OF THE INVENTION

The present invention relates to a recombinant or isolated integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , the subunit  $\alpha 10$  thereof, homologues and fragments of said integrin and of said subunit  $\alpha 10$  having similar biological activity, processes of producing the same, polynucleotides and oligonucleotides encoding the same, vectors and cells comprising the same, binding entities binding specifically to the same, and the use of the same.

#### BACKGROUND OF THE INVENTION

10

The integrins are a large family of transmembrane glycoproteins that mediate cell-cell and cell-matrix 15 interactions (1-5). All known members of this superfamily are non-covalently associated heterodimers composed of an  $\alpha$ - and a  $\beta$ -subunit. At present, 8  $\beta$ -subunits ( $\beta$ 1- $\beta$ 8) (6) and 16  $\alpha$ -subunits ( $\alpha$ 1- $\alpha$ 9,  $\alpha$ v,  $\alpha$ M,  $\alpha$ L,  $\alpha$ X,  $\alpha$ IIb,  $\alpha$ E and 20  $\alpha D$ ) have been characterized (6-21), and these subunits associate to generate more than 20 different integrins. The  $\beta$ 1-subunit has been shown to associate with ten different  $\alpha$ -subunits,  $\alpha 1-\alpha 9$  and  $\alpha v$ , and to mediate interactions with extracellular matrix proteins such as colla-25 gens, laminins and fibronectin. The major collagen binding integrins are  $\alpha 1\beta 1$  and  $\alpha 2\beta 1$  (22-25). The integrins  $\alpha 3\beta 1$  and  $\alpha 9\beta 1$  have also been reported to interact with collagen (26,27) although this interaction is not well understood (28). The extracellular N-terminal regions of the  $\alpha$  and  $\beta$  integrin subunits are important in the bind-30 ing of ligands (29,30). The N-terminal region of the α-subunits is composed of a seven-fold repeated sequence (12,31) containing FG and GAP consensus sequences. The repeats are predicted to fold into a  $\beta$ -propeller domain

30

(32) with the last three or four repeats containing putative divalent cation binding sites. The  $\alpha$ -integrin subunits  $\alpha 1$ ,  $\alpha 2$ ,  $\alpha D$ ,  $\alpha E$ ,  $\alpha L$ ,  $\alpha M$  and  $\alpha X$  contain a ~200 amino acid inserted domain, the I-domain (A-domain), which shows similarity to sequences in von Willebrand factor, cartilage matrix protein and complement factors C2 and B (33,34). The I-domain is localized between the second and third FG-GAP repeats, it contains a metal ion-dependent adhesion site (MIDAS) and it is involved in binding of ligands (35-38).

Chondrocytes, the only type of cells in cartilage, express a number of different integrins including  $\alpha 1\beta 1$ ,  $\alpha 2\beta 1$ ,  $\alpha 3\beta 1$ ,  $\alpha 5\beta 1$ ,  $\alpha 6\beta 1$ ,  $\alpha \nu \beta 3$ , and  $\alpha \nu \beta 5$  (39-41). It has been shown that  $\alpha 1\beta 1$  and  $\alpha 2\beta 1$  mediate chondrocyte interactions with collagen type II (25) which is one of the major components in cartilage. It has also been shown that  $\alpha 2\beta 1$  is a receptor for the cartilage matrix protein chondroadherin (42).

### 20 SUMMARY OF THE INVENTION

The present invention relates to a novel collagen type II binding integrin, comprising a subunit  $\alpha 10$  in association with a subunit  $\beta$ , especially subunit  $\beta 1$ , but also other  $\beta$ -subunits may be contemplated. In preferred embodiments, this integrin has been isolated from human or bovine articular chondrocytes, and human chondrosarcoma cells.

The invention also encompasses integrin homologues of said integrin, isolated from other species, such as bovine integrin heterodimer comprising a subunit  $\alpha 10$  in association with a subunit  $\beta$ , preferably  $\beta 1$ , as well as homologues isolated from other types of human cells or from cells originating from other species.

The present invention relates in particular to a recombinant or isolated integrin subunit  $\alpha 10$  comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, and homologues and or fragments thereof having the

25

35

same biological activity.

The invention further relates to a process of producing a recombinant integrin subunit  $\alpha 10$  comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or homologues or fragments thereof having similar biological activity, which process comprises the steps of

- a) isolating a polynucleotide comprising a nucleotide sequence coding for a integrin subunit  $\alpha 10$ , or homologues or fragments thereof having similar biological activity,
- b) constructing an expression vector comprising the isolated polynucleotide,
- c) transforming a host cell with said expression vector,
- d) culturing said transformed host cell in a culture medium under conditions suitable for expression of integrin subunit  $\alpha 10$ , or homologues or fragments thereof having similar biological activity, in said transformed host cell, and, optionally,
- 20 e) isolating the integrin subunit  $\alpha 10$ , or homologues or fragments thereof having the same biological activity, from said transformed host cell or said culture medium.

The integrin subunit  $\alpha 10$ , or homologues or fragments thereof having the same biological activity, can also be provided by isolation from a cell in which they are naturally present.

The invention also relates to an isolated polynucleotide comprising a nucleotide coding for a integrin subunit  $\alpha 10$ , or homologues or fragments thereof having similar biological activity, which polynucleotide comprises the nucleotide sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or parts thereof.

The invention further relates to an isolated polynucleotide or oligonucleotide which hybridises to a DNA or RNA encoding an integrin subunit  $\alpha 10$ , having the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or homologues or fragments thereof, wherein said polyoligo-

25

30

nucleotide or oligonucleotide fails to hybridise to a DNA or RNA encoding the integrin subunit  $\alpha 1$ .

The invention relates in a further aspect to vectors comprising the above polynucleotides, and to cells containing said vectors and cells that have polynucleotides or oligonycleotides as shown in SEQ ID No. 1 or 2 integrated in their genome.

The invention also relates to binding entities having the capability of binding specifically to the integrin subunit  $\alpha 10$  or to homologues or fragments thereof, such as proteins, peptides, carbohydrates, lipids, natural ligands, polyclonal antibodies or monoclonal antibodies.

In a further aspect, the invention relates to a recombinant or isolated integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , in which the subunit  $\alpha 10$  comprises the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or homologues or fragments thereof having similar biological activity.

In a preferred embodiment thereof, the subunit  $\beta$  is  $\beta$ 1.

The invention also relates to a process of producing a recombinant integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , in which the subunit  $\alpha 10$  comprises the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, which process comprises the steps of

- a) isolating one polynucleotide comprising a nucleotide sequence coding for a subunit  $\alpha 10$  of an integrin heterodimer and, optionally, another polynucleotide comprising a nucleotide sequence coding for a subunit  $\beta$  of an integrin heterodimer, or for homologues or fragments thereof having similar biological activity,
- b) constructing an expression vector comprising said isolated polynucleotide coding for said subunit  $\alpha 10$  in combination with an expression vector comprising said isolated nucleotide coding for said subunit  $\beta$ ,

25

30

- c) transforming a host cell with said expression vectors.
- d) culturing said transformed host cell in a culture medium under conditions suitable for expression of an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or homologues or fragments thereof similar biological activity, in said transformed host cell, and, optionally,
- e) isolating the integrin heterodimer comprising a 10 subunit  $\alpha 10$  and a subunit  $\beta$ , or homologues or fragments thereof having the same biological activity, from said transformed host cell or said culture medium.

The integrin heterodimer, or homologues or fragments thereof having similar biological activity, can also be provided by isolation from a cell in which they are naturally present.

The invention further relates to a cell containing a first vector, said first vector comprising a polynucleotide coding for a subunit  $\alpha 10$  of an integrin heterodimer, 20 or for homologues or parts thereof having similar biological activity, which polynucleotide comprises the nucleotide sequence shown in SEQ ID No. 1 or SEQ ID No. 2 or parts thereof, and, optionally, a second vector, said second vector comprising a polynucleotide coding for a subunit  $\beta$  of an integrin heterodimer, or for homologues or fragments thereof.

In still another aspect, the invention relates to binding entities having the capability of binding specifically to the integrin heterodimer comprising a subunit  $\alpha$ 10 and a subunit  $\beta$ , or to homologues or fragments thereof having similar biological activity, preferably wherein the subunit  $\beta$  is  $\beta$ 1. Preferred binding entities are proteins, peptides, carbohydrates, lipids, natural ligands, polyclonal antibodies and monoclonal antibodies.

35 In a further aspect, the invention relates to a fragment of the integrin subunit  $\alpha 10$ , which fragment is a peptide chosen from the group comprising peptides of

10

15

20

25

30

35

the cytoplasmic domain, the I-domain and the spliced domain.

In one embodiment, said fragment is a peptide comprising the amino acid sequence KLGFFAHKKIPEEEKREEKLEQ.

In another embodiment, said fragment comprises the amino acid sequence from about amino acid no. 952 to about amino acid no. 986 of SEO ID No. 1.

In a further embodiment, said fragment comprises the amino acid sequence from about amino acid No. 140 to about amino acid No. 337 in SEQ ID No. 1.

Another embodiment of the invention relates to a polynucleotide or oligonucleotide coding for a fragment of the human integrin subunit  $\alpha 10$ . In one embodiment this polynucleotide of oligonucleotide codes for a fragment which is a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the spliced domain. In further embodiments the polynucleotide or oligonucleotide codes for the fragments defined above.

The invention also relates to binding entities having the capability of binding specifically to a fragment of the integrin subunit  $\alpha 10$  as defined above.

The invention also relates to a process of using an integrin subunit  $\alpha 10$  comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or an integrin heterodimer comprising said subunit  $\alpha 10$  and a subunit  $\beta$ , or a homologue or fragment of said integrin or subunit having similar biological activity, as a marker or target molecule of cells or tissues expressing said integrin subunit  $\alpha 10$ , which cells or tissues are of animal including human origin.

In an embodiment of this process the fragment is a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the spliced domain.

In further embodiments of said process the fragment is a peptide comprising the amino acid sequence KLGFFAHKKIPEEEKREEKLEQ, or a fragment comprising the amino acid sequence from about amino acid No. 952 to

15

20

25

30

about amino acid No. 986 of SEQ ID No. 1, or a fragment comprising the amino acid sequence from about amino acid no. 140 to about amino acid no. 337 of SEQ ID no. 1.

The subunit  $\beta$  is preferably  $\beta 1$ . The cells are preferably chosen from the group comprising chondrocytes, smooth muscle cells, endothelial cells, osteoblasts and fibroblasts.

Said process may be used during pathological conditions involving said subunit  $\alpha 10$ , such as pathological conditions comprising damage of cartilage, or comprising trauma, rheumatoid arthritis and osteoarthritis.

Said process may be used for detecting the formation of cartilage during embryonal development, or for detecting physiological or therapeutic reparation of cartilage.

Said process may also be used for selection and analysis, or for sorting, isolating or purification of chondrocytes.

A further embodiment of said process is a process for detecting regeneration of cartilage or chondrocytes during transplantation of cartilage or chondrocytes.

A still further embodiment of said process is a process for in vitro studies of differentiation of chondrocytes.

The invention also comprises a process of using binding entities having the capability of binding specifically to an integrin subunit  $\alpha 10$  comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or an integrin heterodimer comprising said subunit  $\alpha 10$  and a subunit  $\beta$ , or to homologues or fragments thereof having similar biological activity, as markers or target molecules of cells or tissues expressing said integrin subunit  $\alpha 10$ , which cells or tissues are of animal including human origin.

The fragment in said process may be a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the spliced domain. In preferred embodiments said fragment is a peptide comprising the

Я

amino acid sequence KLGFFAHKKIPEEEKREEKLEQ, or a fragment comprising the amino acid sequence from about amino acid No. 952 to about amino acid No. 986 of SEQ ID No. 1, or a fragment comprising the amino acid sequence from about amino acid No. 140 to about amino acid no. 337 of SEQ ID No. 1.

The process may also be used for detecting the presence of an integrin subunit  $\alpha 10$  comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or of an integrin heterodimer comprising said subunit  $\alpha 10$  and a subunit  $\beta$ , or of homologues or fragments thereof having similar biological activity.

In a further embodiment said process is a process for determining the differentiation-state of cells during embryonic development, angiogenesis, or development of cancer.

15

In a still further embodiment this process is a process for detecting the presence of an integrin subunit  $\alpha$ 10, or of a homologue or fragment of said integrin 20 subunit having similar biological activity, on cells, whereby a polynucleotide or oligonucleotide chosen from the group comprising a polynucleotide or oligonucleotide chosen from the nucelotide sequence shown in SEQ ID No. 1 is used as a marker under hybridisation conditions where-25 in said polynucleotide or oligonucleotide fails to hybridise to a DNA or RNA encoding an integrin subunit al. Said cells may be chosen from the group comprising chondrocytes, smooth muscle cells, endothelial cells, osteoblasts and fibroblasts. Said integrin fragment may be a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the spliced domain, such as a peptide comprising the amino acid sequence KLGFFAHKKIPEEEKREEKLEQ, or a fragment comprising the amino acid sequence from about amino acid no. 952 to about amino acid no. 986 of SEQ ID No. 1, or a fragment 35 comprising the amino acid sequence from about amino acid No. 140 to about amino acid no. 337 of SEO ID No. 1.

In a still further embodiment the process is a process for determining the differentiation-state of cells during development, in pathological conditions, in tissue regeneration or in therapeutic and physiological reparation of cartilage. The pathological conditions may be any pathological conditions involving the integrin subunit  $\alpha 10$ , such as rheumatoid arthritis, osteoarthrosis or cancer. The cells may be chosen from the group comprising chondrocytes, smooth muscle cells, endothelial cells, osteoblasts and fibroblasts.

The invention also relates to a process for determining the differentiation-state of cells during development, in pathological conditions, in tissue regeneration and in therapeutic and physiological reparation of car-15 tilage, whereby a polynucleotide or oligonucleotide chosen from the nucleotide sequence shown in SEQ ID No. 1 is used as a marker under hybridisation conditions wherein said polynucleotide or oligonucleotide fails to hybridise to a DNA or RNA encoding an integrin subunit α1. Embodiments of this aspect comprise a process, where-20 by said polynucleotide or oligonucleotide is a polynucleotide or oligonucleotide coding for a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the spliced domain, 25 such as a polynucleotide or oligonucleotide coding for a peptide comprising the amino acid sequence KLGFFAHKKIPEEEKREEKLEQ, or comprising the amino acid sequence from about amino acid No. 952 to about amino acid no. 986 of SEQ ID No. 1, or the amino acid sequence 30 from about amino acid No. 140 to about amino acid No. 337 of SEQ ID No. 1. Said pathological conditions may be any pathological conditions involving the integrin subunit α10, such as rheumatoid arthritis, osteoarthrosis or cancer, or atherosclerosis or inflammation. Said cells 35 may be chosen from the group comprising chondrocytes, smooth muscle cells, endothelial cells, osteoblasts and fibroblasts.

15

20

25

30

In a further aspect the invention relates to a pharmaceutical composition comprising as an active ingredient a pharmaceutical agent or an antibody which is capable of using an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit  $\alpha 10$  having similar biological activity, as a target molecule. An embodiment of said pharmaceutical composition is intended for use in stimulating, inhibiting or blocking the formation of cartilage, bone or blood vessels. A further embodiment comprises a pharmaceutical composition for use in preventing adhesion between tendon/ligaments and the surrounding tissue after infection, inflammation and after surgical intervention where adhesion impairs the function of the tissue.

The invention is also related to a vaccine comprising as an active ingredient an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit  $\alpha 10$ , or DNA or RNA coding for said integrin subunit  $\alpha 10$ .

A further aspect of the invention is related to the use of the integrin subunit  $\alpha 10$  as defined above as a marker or target in transplantation of cartilage or chondrocytes.

A still further aspect of the invention is related to a method of using binding entities having the capability of binding specifically to an integrin subunit  $\alpha 10$  comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or an integrin heterodimer comprising said subunit  $\alpha 10$  and a subunit  $\beta$ , or to homologues or fragments thereof having similar biological activity, for promoting adhesion of chondrocytes and/or osteoblasts to surfaces of implants to stimulate osseointegration.

35 The invention is also related to the use of an integrin subunit  $\alpha 10$  or an integrin heterodimer comprising said subunit  $\alpha 10$  and a subunit  $\beta$  as a target for anti-

15

20

25

30

35

adhesive drugs or molecules in tendon, ligament, skeletal muscle or other tissues where adhesion impairs the function of the tissue.

The invention also relates to a method of stimulating, inhibiting or blocking the formation of cartilage or bone, comprising administration to a subject a suitable amount of a pharmaceutical agent or an antibody which is capable of using an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit  $\alpha 10$  having similar biological activity, as a target molecule.

In another embodiment the invention is related to a method of preventing adhesion between tendon/ligaments and the surrounding tissue after infection, inflammation and after surgical intervention where adhesion impairs the function of the tissue, comprising administration to a subject a suitable amount of a pharmaceutical agent or an antibody which is capable of using a integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit  $\alpha 10$  having similar biological activity, as a target molecule.

The invention also relates to a method of stimulating extracellular matrix synthesis and repair by activation or blockage of an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or of the subunit  $\alpha 10$  thereof, or of a homologue or fragment of said integrin or subunit  $\alpha 10$  having similar biological activity.

In a further aspect the invention relates to a method of in vitro detecting the presence of integrin binding entities, comprising interaction of an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit, with a sample, thereby causing said integrin, subunit  $\alpha 10$ , or homologue or fragment thereof having similar biological activity, to modulate

the binding to its natural ligand or other integrin binding proteins present in said sample.

The invention also relates to a method of in vitro studying consequences of the interaction of a human heterodimer integrin comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit, with an integrin binding entity and thereby initiate a cellular reaction. Said consequences may be measured as alterations in cellular functions.

A still further aspect of the inventions relates to a method of using DNA or RNA encoding an integrin subunit  $\alpha 10$  or homologues or fragments thereof as a molecular target. In an embodiment of this aspect, a polynucleotide or oligonucleotide hybridises to the DNA or RNA encoding an integrin subunit  $\alpha 10$  or homologues or fragments thereof, whereby said polynucleotide or oligonucleotide fails to hybridise to a DNA or RNA encoding en integrin subunit  $\alpha 1$ .

The invention also relates to a method of using a human heterodimer integrin comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit, or a DNA or RNA encoding an integrin subunit  $\alpha 10$  or homologues or fragments thereof, as a marker or target molecule during angiogenesis.

BRIEF DESCRIPTION OF THE FIGURES

Fig.1 Affinity purification of the  $\alpha 10$  integrin subunit on collagen type II-Sepharose.

30 Fig. 2. Amino acid sequences of peptides from the bovine  $\alpha 10$  integrin subunit.

Fig. 3a. Affinitypurification and immunoprecipitation of the integrin subunit  $\alpha 10$  from bovine chondrocytes.

Fig. 3b. Affinitypurification and immunoprecipitation of the integrin subunit  $\alpha 10$  from human chondrocytes.

WO 99/51639

10

25

35

Fig. 3c. Affinitypurification and immunoprecipitation of the integrin subunit  $\alpha 10$  from human chondrosarcoma cells.

Fig. 4. A 900 bp PCR-fragment corresponding to the bovine integrin subunit  $\alpha 10$ 

Fig. 5. Schematic map of the three overlapping  $\alpha 10\,$  clones.

Fig. 6. Nucleotide sequence and deduced amino acid sequence of the human  $\alpha 10$  integrin subunit.

Fig. 7. Northern blot of integrin alo mRNA.

Fig. 8 Immunoprecipitation of the  $\alpha 10$  integrin subunit from human chondrocytes using antibodies against the cytoplasmic domain of  $\alpha 10$  (a). Western blot of the  $\alpha 10$  associated  $\beta$ -chain (b).

15 Fig. 9. Immunostaining of  $\alpha$ 10 integrin in human articular cartilage.

Fig. 10 Immunostaining of  $\alpha 10$  integrin in 3 day mouse limb cartilage.

Fig 11. Immunostaining of  $\alpha$ 10 integrin in 13.5 day 20 mouse embryo.

Fig 12. Hybridisation of  $\alpha 10~\text{mRNA}$  in various human tissues.

Fig. 13 Immunostaining of fascia around tendon (a), skeletal muscle (b) and heart valves (c) in 3 day mouse limb.

Fig. 14. PCR fragments corresponding to  $\alpha 10$  integrin subunit from human chondrocytes, human endothelial cells, human fibroblasts and rat tendon.

Fig 15. Partial genomic nucleotide sequence of the 30 human integrin subunit  $\alpha 10\,.$ 

Fig 16. Upregulation of  $\alpha 10$  integrin subunit in chondrocytes cultured in alginate.

Fig 17. Immunoprecipitation of the  $\alpha 10$  integrin subunit from human smooth muscle cells

DETAILED DESCRIPTION OF THE INVENTION

The present invention demonstrate that human and

14

bovine chondrocytes express a novel, collagen type II-binding integrin in the  $\beta$ 1-family. An earlier study presented some evidence for that human chondrosarcoma cells also express this integrin (25). Immunoprecipitation experiments using antibodies against the integrin subunit  $\beta 1$  revealed that this novel  $\alpha$ -integrin subunit had an apparent molecular weight  $(M_r)$  of approximately 160 kDa under reducing conditions, and was slightly larger than the  $\alpha 2$  integrin subunit. To isolate this α-subunit collagen type II-binding proteins were affinity 10 purified from bovine chondrocytes. The chondrocyte lysate was first applied to a fibronectin-Sepharose precolumn and the flow through was then applied to a collagen type II-Sepharose column. A protein with Mr of approximately 160 kD was specifically eluted with EDTA from the colla-15 gen column but not from the fibronectin column. The  $M_{\rm r}$  of this protein corresponded with the  $M_{\mbox{\scriptsize r}}$  of the unidentified β1-related integrin subunit. The 160 kD protein band was excised from the SDS-PAGE gel, digested with trypsin and 20 the amino acid sequences of the isolated peptides were analysed.

Primers corresponding to isolated peptides amplified a 900 bp PCR-fragment from bovine cDNA which was cloned, sequenced and used for screening of a human articular chondrocyte  $\lambda \mathrm{ZapII}$  cDNA library to obtain the human integrin  $\alpha$ -subunit homologue. Two overlapping clones, hcl and hc2 were isolated, subcloned and sequenced. These clones contained 2/3 of the nucleotide sequence including the 3' end of the cDNA. A third clone which contained the 5'end of the  $\alpha$ 10 cDNA, was obtained using the RACE technique. Sequence analysis of the 160 kD protein sequence showed that it was a member of the integrin  $\alpha$ -subunit family and the protein was named  $\alpha$ 10.

25

35

The deduced amino acid sequence of  $\alpha 10$  was found to share the general structure of the integrin  $\alpha$ -subunits described in previously published reports (6-21). The large extracellular N-terminal part of  $\alpha 10$  contains a

seven-fold repeated sequence which was recently predicted to fold into a  $\beta$ -propeller domain (32). The integrin subunit  $\alpha 10$  contains three putative divalent cation-binding sites (DxD/NxD/NxxxD) (53), a single spanning transmembrane domain and a short cytoplasmic domain. In contrast to most  $\alpha$ -integrin subunits the cytoplasmic domain of  $\alpha$ 10 does not contain the conserved sequence KxGFF (R/K) R. The predicted amino acid sequence in  $\alpha 10$  is KLGFFAH. Several reports indicate that the integrin cytoplasmic 10 domains are crucial in signal transduction (54) and that membrane-proximal regions of both  $\alpha$ - and  $\beta$ -integrin cytoplasmic domains are involved in modulating conformation and affinity state of integrins (55-57). It is suggested that the GFFKR motif in  $\alpha$ -chains are important for association of integrin subunits and for transport of the integrin to the plasma membrane (58). The KxGFFKR domain has been shown to interact with the intracellular protein calreticulin (59) and interestingly, calreticulin-null. embryonic stem cells are deficient in integrin-mediated 20 cell adhesion (60). It is therefor possible that the sequence KLGFFAH in all have a key function in regulating the affinity between  $\alpha 10\beta 1$  and matrix proteins.

Integrin  $\alpha$  subunits are known to share an overall identity of 20-40% (61). Sequence analysis showed that 25 the  $\alpha 10$  subunit is most closely related to the I-domain containing  $\alpha$ -subunits with the highest identity to  $\alpha 1$ (37%) and  $\alpha$ 2 (35%). The integrins  $\alpha$ 1 $\beta$ 1 and  $\alpha$ 2 $\beta$ 1 are known receptors for both collagens and laminins (24;62;63) and we have also recently demonstrated that  $\alpha 2\beta 1$  interacts with the cartilage matrix protein chondroadherin (42). Since  $\alpha 10\beta 1$  was isolated on a collagen type II-Sepharose, we know that collagen type II is a ligand for  $\alpha 10\beta 1$ . We have also shown by affinity purification experiments that  $\alpha$ 10 $\beta$ 1 interacts with collagen type I but it remains to be 35 seen whether laminin or chondroadherin are also ligands for this integrin.

16

The  $\alpha 10$  associated  $\beta$ -chain migrated as the  $\beta 1$  integrin subunit both under reducing and non-reducing conditions. To verify that the  $\alpha 10$  associated  $\beta$ -chain indeed is  $\beta 1$ , chondrocyte lysates were immunoprecipitated with antibodies against  $\alpha 10$  or  $\beta 1$  followed by Western blot using antibodies against the  $\beta 1$ -subunit. These results clearly demonstrated that  $\alpha 10$  is a member of the  $\beta 1$ -integrin family. However, the possibility that  $\alpha 10$  combine also with other  $\beta$ -chains can not be excluded.

10

15

20

25

30

35

A polyclonal peptide antibody raised against the cytoplasmic domain of  $\alpha 10$  precipitated two protein bands with  $M_r$  of approximately 160 kD ( $\alpha$ 10) and 125 kD ( $\beta$ 1) under reducing conditions. Immunohistochemistry using the  $\alpha$ 10-antibody showed staining of the chondrocytes in tissue sections of human articular cartilage. The antibody staining was clearly specific since preincubation of the antibody with the  $\alpha$ 10-peptide completely abolished the staining. Immunohistochemical staining of mouse limb sections from embryonic tissue demonstrated that  $\alpha 10$  is upregulated during condensation of the mesenchyme. This indicate that the integrin subunit  $\alpha 10$  is important during the formation of cartilage. In 3 day old mice  $\alpha 10$ was found to be the dominating collagen binding integrin subunit which point to that  $\alpha 10$  has a key function in maintaining normal cartilage functions.

Expression studies on the protein and mRNA level show that the distribution of  $\alpha 10$  is rather restrictive. Immunohistochemistry analyses have shown that  $\alpha 10$  integrin subunit is mainly expressed in cartilage but it is also found in perichondrium, periosteum, ossification groove of Ranvier, in fascia surrounding tendon and skeletal muscle and in the tendon-like structures in the heart valves. This distribution point to that  $\alpha 10$  integrin subunit is present also on fibroblasts and osteoblasts. PCR amplification of cDNA from different cell types revealed the presence of an alternatively spliced  $\alpha 10$  integrin subunit. This spliced  $\alpha 10$  was domi-

nating in fibroblasts which suggests that α10 in fibroblasts may have a different function compared to alo present on chondrocytes.

Expression of the integrin subunit  $\alpha 10$  was found to decrease when chondrocytes were cultured in monolayer. In contrast, the expression of  $\alpha 10$  was found to increase when the cells were cultured in alginate beads. Since the latter culturing model is known to preserve the phenotype of chondrocytes the results suggest that  $\alpha 10$  can function as marker for a differentiated chondrocyte.

Adhesion between tendon/ligaments and the surrounding tissue is a well-known problem after infection, injury and after surgical intervention. Adhesion between tendon and tendon sheets impairs the gliding function and cause considerable problems especially during healing of tendons in e.g. the hand and fingers leading to functional incapacity. The localisation of the  $\alpha 10$  integrin subunit in the fascia of tendon and skeletal muscle makes α10 a possible target for drugs and molecules with anti-20 adhesive properties that could prevent impairment of the function of tendon/ligament. The integrin subunit  $\alpha 10$  can also be a target for drugs or molecules with anti-adhesive properties in other tissues where adhesion is a problem.

25

10

15

# **EXAMPLES**

#### Example 1

Affinity purification of the α10integrin subunit on 30 collagen type II-Sepharose.

Materials and Methods

Bovine chondrocytes, human chondrocytes or human chondrosarcoma cells were isolated as described earlier [Holmvall et al, Exp Cell Res, 221, 496-503 (1995), Camper et al, JBC, 273, 20383-20389 (1998)]. A Triton X-100 lysate of bovine chondrocytes was applied to a fibronectin-Sepharose precolumn followed by a collagen

18

type II-Sepharose column and the integrin subunit  $\alpha10$  was eluted from the collagen type II-column by EDTA (Camper et al, JBC, 273, 20383-20389 (1998). The eluted proteins were precipitated by methanol/chloroform, separated by SDS-PAGE under reducing conditions and stained with Coomassie blue. (Camper et al, JBC, 273, 20383-20389 (1998). Peptides from the  $\alpha10$  protein band were isolated by in-gel digestion with a trypsin and phase liquid chromatography and sequenced by Edman degradation (Camper et al, JBC, 273, 20383-20389 (1998). Results

Fig 1 shows EDTA-eluted proteins from the fibronectin-Sepharose (A), flow-through from the collagen type II-Sepharose column (B) and EDTA-eluted proteins from the collagen type II-Sepharose (C). The  $\alpha 10$  integrin subunit (160 kDa) which was specifically eluted from the collagen type II column is indicated with an arrow. Figure 2 shows the amino acid sequences of six peptides that were isolated from the bovine integrin subunit  $\alpha 10$ . Figures 3 a, b, and c show that the  $\alpha 10$  integrin subunit is present on bovine chondrocytes (3a), human chondrocytes (3b) and human chondrosarcoma cells (3c). The affinity for collagen type II, the coprecipitation with  $\beta 1$ -integrin subunit and the molecular weight of 160 kDa under reducing conditions identify the  $\alpha 10$  integrin subunit on the different

cells. These results show that  $\alpha 10$  can be isolated from

## Example 2

10

15

25

30 Amplification of PCR fragment corresponding to bovine  $\alpha 10$  integrin subunit. Materials and methods

chondrocytes and from chondrosarcoma cells.

The degenerate primers GAY AAY ACI GCI CAR AC (DNTAQT, forward) and TIA TIS WRT GRT GIG GYT (EPHHSI, reverse) were used in PCR (Camper et al, JBC, 273, 20383-20389 (1998) to amplify the nucleotide sequence corresponding to the bovine peptide 1 (Figure 2). A 900 bp

PCR-fragment was then amplified from bovine cDNA using an internal specific primer TCA GCC TAC ATT CAG TAT (SAYIQY, forward) corresponding to the cloned nucleotide sequence of peptide 1 together with the degenerate primer ICK RTC CCA RTG ICC IGG (PGHWDR, reverse) corresponding to the bovine peptide 2 (Figure2). Mixed bases were used in positions that were twofold degenerate and inosines were used in positions that are three- or fourfold degenerate. mRNA isolation and cDNA synthesis was done as earlier described (Camper et al, JBC, 273, 20383-20389 (1998)). The purified fragment was cloned, purified and sequenced as earlier described (Camper et al, JBC, 273, 20383-20389 (1998)).

Results

The nucleotide sequence of peptide 1 (Figure 2)
was obtained by PCR-amplification, cloning and sequencing of bovine cDNA. From this nucleotide sequence an
exact primer was designed and applied in PCR-amplification with degenerate primers corresponding to peptides
20 2-6 (Figure 2). Primers corresponding to peptides 1
and 2 amplified a 900 bp PCR-fragment from bovine cDNA
(Figure 4).

## Example 3

35

25 Cloning and sequence analysis of the human  $\alpha 10$  integrin subunit

Material and methods

The cloned 900bp PCR-fragment, corresponding to bovine  $\alpha$ 10-integrin, was digoxigenin-labelled according to the DIG DNA labelling kit (Boehringer Mannheim) and used as a probe for screening of a human articular chondrocyte  $\lambda$ ZapII cDNA library (provided by Michael Bayliss, The Royal Veterinary Basic Sciences, London, UK) (52). Positive clones containing the pBluescript SK+ plasmid with the cDNA insert were rescued from the ZAP vector by in vivo excision as described in the ZAP-cDNA® synthesis kit (Stratagene). Selected plasmids were purified and

20

sequenced as described earlier (Camper et al, JBC, 273, 20383-20389 (1998)) using T3, T7 and internal specific primers. To obtain cDNA that encoded the 5' end of α10 we designed the primer AAC TCG TCT TCC AGT GCC ATT CGT GGG (reverse; residue 1254-1280 in α10 cDNA) and used it for rapid amplification of the cDNA 5' end (RACE) as described in the Marathon™ cDNA Amplification kit (Clontech INC., Palo Alto, CA).

Results

Two overlapping clones, hcl and hc2 (Figure 5), were 10 isolated, subcloned and sequenced. These clones contained 2/3 of the nucleotide sequence including the 3' end of the cDNA. A third clone (racel; Figure 5), which contained the 5'end of the  $\alpha$ 10 cDNA, was obtained using the RACE technique. From these three overlapping clones of 15 alo cDNA, 3884 nucleotides were sequenced The nucleotide sequence and deduced amino acid sequence is shown in Figure 6. The sequence contains a 3504-nucleotide open reading frame that is predicted to encode a 1167 amino acid mature protein. The signal peptide cleavage site is 20 marked with an arrow, human homologues to bovine peptide sequences are underlined and the I-domain is boxed. Metal ion binding sites are indicated with a broken underline, potential N-glycosylation sites are indicated by an 25 asterisk and the putative transmembrane domain is double underlined. The normally conserved cytoplasmic sequence is indicated by a dot and dashed broken underline.

Sequence analysis demonstrate that  $\alpha 10$  is a member of the integrin  $\alpha$ -subunit family.

Example 4

30

Identification of a clone containing a splice variant of  $\alpha 10$ 

One clone which was isolated from the human chon- drocyte library (see Example 3) contained a sequence that was identical to the sequence of  $\alpha 10$  integrin subunit except that the nucleotides between nt positions

21

2942 and 3055 were deleted. The splice variant of  $\alpha 10$  was verified in PCR experiment using primers flanking the splice region (see figure 14).

## 5 Example 5

30

35

Identification of  $\alpha 10$  integrin subunit by Northern blot

Material and methods

Bovine chondrocyte mRNA was purified using a QuickPrep®Micro mRNA Purification Kit (Pharmacia Biotech, 10 Uppsala, Sweden), separated on a 1% agarose-formaldehyde gel, transferred to nylon membranes and immobilised by UVcrosslinking. cDNA-probes were 32P-labelled with Random Primed DNA Labeling Kit (Boehringer Mannheim). Filters 15 were prehybridised for 2-4 hours at 42°C in 5x SSE, 5x Denharts solution, 0.1 % SDS, 50 μg/ml salmon sperm DNA and 50% formamide and then hybridised over night at 42 °C with the same solution containing the specific probe (0.5-1 x 106 cpm/ml). Specifically bound cDNA-20 probes were analysed using the phosphoimager system (Fuji). Filters were stripped by washing in 0.1% SDS, for 1 hour at  $80^{\circ}$ C prior to re-probing. The  $\alpha$ 10-integrin cDNA-probe was isolated from the racel-containing plasmid using the restriction enzymes BamHI (GIBCO BRL) and NcoI 25 (Boehringer Mannheim). The rat \$1-integrin cDNA probe was a kind gift from Staffan Johansson, Uppsala, Sweden. Results

Northern blot analysis of mRNA from bovine chondrocytes showed that a human  $\alpha 10$  cDNA-probe hybridised with a single mRNA of approximately 5.4 kb (Figure 7). As a comparison, a cDNA-probe corresponding to the integrin subunit  $\alpha 1$  was used. This cDNA-probe hybridised a mRNA-band of approximately 3.5 kb on the same filter. These results show that a cDNA-probe against  $\alpha 10$  can be used to identify the  $\alpha 10$  integrin subunit on the mRNA level.

## Example 6

Preparation of antibodies against the integrin subunit  $\alpha 10$ 

A peptide corresponding to part of the  $\alpha 10$  cytoplasmic domain, Ckkipeeekreekle (see figure 6) was synthesised and conjugated to keyhole limpet hemocyanin (KLH). Rabbits were immunised with the peptide-KLH conjugate to generate antiserum against the integrin subunit  $\alpha 10$ . Antibodies recognising  $\alpha 10$  were affinity purified on an peptide-coupled column (Innovagen AB).

## Example 7

10

Immunoprecipitation of the integrin subunit  $\alpha 10\ \text{from}$  chondrocytes

# 15 Material and methods

Human chondrocytes were 125I-labelled, lyzed with Triton X-100 and immunoprecipitated as earlier described (Holmvall et al, Exp Cell Res, 221, 496-503 (1995), Camper et al, JBC, 273, 20383-20389 (1998)). Triton X-100 lysates of 125I-labeled human chondrocytes were immuno-20 precipitated with polyclonal antibodies against the integrin subunits  $\beta1$ ,  $\alpha1$ ,  $\alpha2$ ,  $\alpha3$  or  $\alpha10$ . The immunoprecipitated proteins were separated by SDS-PAGE (4-12%) under non-reducing conditions and visualised using a phospho-25 imager. Triton X-100 lysates of human chondrocytes immunoprecipitated with  $\alpha 10$  or  $\beta 1$  were separated by SDS-PAGE (8%) under non-reducing conditions and analysed by Western blot using the polyclonal \$1 antibody and chemiluminescent detection as described in Camper et al, JBC, 30 273, 20383-20389 (1998).

#### Results

35

The polyclonal peptide antibody, raised against the cytoplasmic domain of  $\alpha 10$ , precipitated two protein bands with Mr of approximately 160 kD ( $\alpha 10$ ) and 125 kD ( $\beta 1$ ) under reducing conditions. The  $\alpha 10$  associated  $\beta$ -chain migrated as the  $\beta 1$  integrin subunit (Figure 8a). To verify that the  $\alpha 10$  associated  $\beta$ -chain in chondrocytes

indeed is  $\beta 1$ , chondrocyte lysates were immunoprecipitated with antibodies against  $\alpha 10$  orb  $\beta 1$  followed by Western blot using antibodies against the  $\beta 1$ -subunit (Figure 8b). These results clearly demonstrated that  $\alpha 10$  is a member of the  $\beta 1$ -integrin family. However, the results do not exclude the possibility that  $\alpha 10$  can associate with other  $\beta$ -chains in other situations.

## Example 8

30

35

Immunohistochemical staining of the integrin subunit  $\alpha 10$  in human and mouse cartilage Material and methods

Frozen sections of adult cartilage (trochlear groove) obtained during surgery (provided by Anders

Lindahl, Salgrenska Hospital, Gothenburg, Sweden and frozen sections from of 3 day old mouse limb were fixed and prepared for immunohistochemistry as earlier described (Camper et al, JBC, 273, 20383-20389 (1998)). Expression of α10 integrin subunit was analysed using the polyclonal antibody against the cytoplasmic domain as a primary antibody (see Example 6) and a secondary antibody conjugated to peroxidase.

Results

Figures 9 show immunostaining of human adult articu-25 lar cartilage.

The  $\alpha 10$ -antibody recognising the cytoplasmic domain of  $\alpha 10$  stained the chondrocytes in tissue sections of human articular cartilage (A). The staining was depleted when the antibody was preincubated with the  $\alpha 10$ - peptide (B). A control antibody recognising the  $\alpha 9$  integrin subunit did not bind to the chondrocyte (C).

Figures 10 shows that the  $\alpha 10$  antibody stain the majority of chondrocytes in the growing bone anlage (a and b). The  $\alpha 10$  antibody also recognised cells in the ossification groove of Ranvier (b), especially the osteoblast in the bone bark which are lining the cartilage in the metaphys are highly positive for  $\alpha 10$ . The

cells in the ossification groove of Ranvier are believed to be important for the growth in diameter of the bone. The integrin subunit  $\alpha 10$  is also highly expressed in perichondrium and periosteum. Cell in these tissues are likely important in the repair of the cartilage tissue. The described localisation of the integrin subunit  $\alpha 10$  suggest that this integrin is important for the function of the cartilage tissue.

## 10 Example 9

15

20

Immunohistochemical staining of the integrin subunit  $\alpha 10$  during mouse development  $$\operatorname{\textsc{Material}}$$  and methods

Frozen sections from mouse embryos (13.5 days) were investigated for expression of  $\alpha 10$  by immunhistochemistry as described in Camper et al, JBC, 273, 20383-20389 (1998). Expression of  $\alpha 10$  integrin subunit was analysed using the polyclonal antibody against the cytoplasmic domain as a primary antibody (see Example 6) and a secondary antibody conjugated to peroxidase. The embryo sections were also investigated for expression of integrin subunit  $\alpha 1$  (monoclonal antibody from Pharmingen) and collagen type II (monoclonal antibody, kind gift from Dr John Mo, Lund University, Sweden).

#### 25 Results

Figure 11 show that α10 integrin subunit is unregulated in the limb when the mesenchymal cells undergo condensation to form cartilage (a). Especially the edge of the newly formed cartilage has high expression of α10.

30 The formation of cartilage is verified by the high expression of the cartilage specific collage type II (b). The control antibody against α1 integrin subunit showed only weak expression on the cartilage (c). In other experiments expression of α10 was found in all cartilage containing tissues in the 3 day old mouse including limbs, ribs and vertebrae. The upregulation of α10 during formation of cartilage suggest that this integrin subunit is

important both in the development of cartilage and bone and in the repair of damaged cartilage tissue.

## Example 10

5

15

mRNA expression of  $\alpha 10$  in tissues other than articular cartilage

Material and methods

Expression of  $\alpha 10$  integrin subunit was examined on the mRNA level in different human tissues. A Northern dot blot with immobilised mRNA from the listed tissues in Figure 12 was hybridised with an  $\alpha 10$  integrin cDNA probe isolated from the race 1-containing plasmid using the restriction enzymes Bam H1 and Nco1. The degree of hybridisation was analysed using a phospho imager. The following symbols denote mRNA level in increasing order: -, +, +++, ++++.

## Results

Analysis of the hybridised mRNA showed that  $\alpha 10$  was expressed in aorta, trachea, spinal cord, heart, lung, and kidney (Figure 12). All other tissues appeared negative for  $\alpha 10$  expression. These results point to a restricted distribution of the  $\alpha 10$  integrin subunit.

# Example 11

Immunohistochemical staining of  $\alpha 10$  in fascia around tendon and skeletal muscle and in tendon structures in heart valves.

## Materials and methods

Frozen sections of adult cartilage (trochlear groove) obtained during surgery (provided by Anders Lindahl, Salgrenska Hospital, Gothenburg, Sweden and frozen sections from of 3 day old mouse limb were fixed and prepared for immunohistochemistry as earlier described (Camper et al, JBC, 273, 20383-20389 (1998)). Expression of α10 integrin subunit was analysed using the polyclonal antibody against the cytoplasmic domain as a pri-

mary antibody (see Example 6) and a secondary antibody conjugated to peroxidase.
Results

As shown in figures 13 expression of  $\alpha 10$  was found in the fascia surrounding tendon (a) and skeletal muscle (b) and in the tendon structures in the heart valves (c). This localisation suggest that  $\alpha 10$  can bind to other matrix molecules in addition to the cartilage specific collagen type II. The localisation of the integrin  $\alpha 10$  on the surface of tendons indicate that  $\alpha 10$  can be involved in unwanted adhesion that often occurs between tendon/ligaments and the surrounding tissue after infection, injury or after surgery.

# 15 Example 12

mRNA expression of  $\alpha 10$  integrin subuhit in chondrocytes, endothelial cells and fibroblasts. Material and methods

Isolation of mRNA, synthesis of cDNA and PCR ampli-20 fication was done as earlier described (Camper et al, JBC, 273, 20383-20389 (1998)). Results

Figure 14 shows PCR amplification of  $\alpha$ 10 cDNA from human articular chondrocytes (lanes A6 and B1), human 25 umbilical vein endothelial cells (lane A2), human fibroblasts (lane A4) and rat tendon (Fig 14b, lane B2). Lanes 1, 3, and 5 in figure 14 A show amplified fragments corresponding to the integrin subunit  $\alpha 2$  in endothelial cells, fibroblasts and chondrocytes, respectively. cDNA-30 primers corresponding to the  $\alpha 10$  sequence positions nt 2919-2943 (forward) and nt 3554-3578 (reverse) (see Figure 6) were used to amplify  $\alpha 10$  cDNA from the different cells. The figure shows that  $\alpha 10$  was amplified in all three cell types. Two fragments of  $\alpha 10$  was amplified which represent the intact form of  $\alpha 10$  (larger fragment) and a splice variant (smaller fragment). The larger fragment was dominating in chondrocytes while the smaller fragment was more pronounced in tendon (B2).

## Example 13

Construction of  $\alpha 10$  mammalian expression vector. 5 The full length protein coding sequence of  $\alpha 10$  (combined from 3 clones, see figure 6) was inserted into the mammalian expression vector, pcDNA3.1/Zeo (Invitrogen). The vector contains SV40 promoter and Zeosin selection sequence. The all containing expression vector was trans-10 fected into cells that express the  $\beta1$ -integrin subunit but lack expression of the  $\alpha 10$  subunit. Expression of the α10 integrin subunit on the cell surface can be analysed by immunoprecipitation and/or flow cytometry using antibodies specific for  $\alpha 10$ . The ligand binding capacity and 15 the function of the inserted all integrin subunit can be demonstrated in cell adhesion experiment and in signalling experiments.

## 20 Example 14

25

Construction of mammalian expression vector containing a splice variant of  $\alpha 10$ .

The full length protein coding sequence of the splice variant of  $\alpha 10$  (nt 2942-nt3055 deleted) was inserted into the mammalian expression vector pcDNA3 (see Example 13). Expression and function of the splice variant can be analysed as described in example 13 and compared with the intact  $\alpha 10$  integrin subunit.

## 30 Example 15

Partial isolation and characterisation of the  $\alpha 10$  integrin genomic DNA \$Material\$ and methods

Human α10 cDNA, isolated from the racel-containing plasmid using the restriction enzymes BamHI (GIBCO BRL) and NcoI (Boehringer Mannheim), was <sup>32</sup>P-labelled and used as a probe for screening of a mouse 129 cosmid library

(provided by Reinhard Fässler, Lund University). Positive clones were isolated and subcloned. Selected plasmids were purified and sequenced as described earlier (Camper et al, JBC, 273, 20383-20389 (1998)) using T3, T7 and internal specific primers. Primers corresponding to mouse genomic DNA were then constructed and used in PCR to amplify and identify the genomic sequence of  $\alpha$ 10 from the cosmid clones.

#### Results

Figure 15 shows 7958 nt of the  $\alpha 10$  gene. This partial genomic DNA sequence of  $\alpha 10$  integrin contains 8 exons, and a Kozak sequence. The mouse genomic  $\alpha 10$  sequence was used to generate a targeting vector for knockout experiments.

15

10

## Example 16

Upregulation of  $\alpha 10$  integrin subunit in chondrocytes cultured in alginate beads Material and methods

Human chondrocytes cultured in monolayer for 2 weeks were detached with trypsin-EDTA and introduced into alginate beads. Chondrocytes cultured in alginate are known to preserve their phenotype while chondrocytes cultured in monolayer are dedifferentiated. After 11 days chondrocytes cultured either in alginate or on monolayer were isolated and surface labelled with  $^{125}\mathrm{I}$ . The  $\alpha10$  integrin subunit was then immunoprecipitated with polyclonal antibodies recognising the cytoplasmic domain of  $\alpha10$  (see Example 6 and Camper et al, JBC, 273, 20383-20389 30 (1998)).

## Results -

35

As shown in figure 16 chondrocytes cultured in alginate beads (lanes 3 and 4) upregulated their protein expression of  $\alpha 10\beta 1$ . This was in contrast to chondrocytes cultured in monolayer (lanes 1 and 2) which had a very low expression of  $\alpha 10\beta 1$ . Immunoprecipitation with ab control antibody is shown in lanes 1 and 3.It is known that

chondrocytes preserve their cartilage specific matrixproduction in alginate cultures but not in monolayer culture which point to that alginate preserve the phenotype of chondrocytes. These results support that  $\alpha 10$  integrin subunit can be used as a marker for differentiated chondrocytes.

#### Example 17

 $\label{eq:continuous} Immunoprecipitation of the $\alpha 10$ integrin subunit from $10$ human smooth muscle cells. $$Material and methods$ 

Human smooth muscle cells were isolated from human aorta. After one week in culture the cells were  $^{125}\text{I}-$  labelled, lysed and immunoprecipitated with antibodies against the integrin subunit  $\beta 1$  (lane 1),  $\alpha 1$  (lane 2),  $\alpha 2$  (lane 3),  $\alpha 10$  (lane 4),  $\alpha 3$  (lane 5), control (lane 6) (Figure 17). The experiment was done as described in Example 7.

#### Results

15

20 The  $\alpha 10$  antibody precipitated two bands from the smooth muscle cells corresponding to the  $\alpha 10$  and the  $\beta 1$  integrin subunit (Fig. 17).

# Example 18

25 Construction of bacterial expression vector containing sequence for  $\alpha 10$  splice region.

A plasmid for intracellular expression in E. coli of the alternatively spliced region (amino acid pos. 952-986, SEQ. ID 1) was constructed as described. The alternatively spliced region were back-translated using the E. coli high frequency codon table, creating a cDNA sequence of 96% identity with the original sequence (SEQ. ID 1 nucleotide pos 2940-3044). Using sequence overlap extension (Horton et al., Biotechniques 8:528, 1990) primer α10pfor (tab. I) and α10prev (tab. I) was used to generate a double stranded fragment encoding the α10 amino acid sequence. This fragment was used as a PCR

template with primers  $\alpha 10 pfor 2$  (tab. I) and  $\alpha 10 prev 2$  (tab. I) in order to generate restriction enzyme site for sub-cloning in a pET vector containing the Z-domain of staphylococcal protein A, creating a fusion of the  $\alpha 10$  spliced region with the amino terminal of the Z-domain with trombin cleavage site residing in-between. The fragment generated in the second PCR reaction is shown (SEQ ID No. 3) also indicating the unique restriction enzymes used for sub-cloning in the expression vector.

10

Table I

al0pfor	5'- GTTCAGAACCTGGGTTGCTACGTTGTTTCCGGTCTGATCATCTCCGC TCTGCTGCCGGCTGT-3'
α10pfor2	5'-GGGGCATATGGTTCAGAACCTGGGTTGCTACGTTG-3'
α10prev	5'- GATAACCTGGGACAAGCTTAGGAAGTAGTTACCACCGTGAGCAACAG CCGGCAGCAGAGCGGA-3'
α10prev2	5'- GGGGGGATCCGCGCGCACCAGGCCGCTGATAACCTGGGACAAGCTT AGGAAGT-3'

## REFERENCES

- 1. Springer, T.A. (1990) Nature 346, 425-434
- 2. Ruoslahti, E. (1991) J.Clin.Invest. 87, 1-5
- 3. Hynes, R.O. (1992) Cell 69, 11-25
- 5 4. Hemler, M.E. (1988) Immunol. Today 9, 109-113
  - 5. Yamada, K.M. (1991) J.Biol.Chem. 266, 12809-12812
  - Palmer, E.L., Ruegg, C., Ferrando, R., Pytela, R., and Sheppard, D. (1993) J.Cell Biol. 123, 1289-1297
- 7. Takada, Y., Elices, M.J., Crouse, C., and Hemler, 10 M.E. (1989) EMBO J. 8, 1361-1368
  - 8. Poncz, M., Eisman, R., Heidenreich, R., Silver, S.M., Vilaire, G., Surrey, S., Schwartz, E., and Bennett, J.S. (1987) J.Biol.Chem. 262, 8476-8482
- Larson, R.S., Corbi, A.L., Berman, L., and Springer,
   T. (1989) J.Cell Biol. 108, 703-712
  - 10. Corbi, A.L., Kishimoto, T.K., Miller, L.J., and Springer, T.A. (1988) J.Biol.Chem. 263, 12403-12411
  - 11. Argraves, W.S., Suzuki, S., Arai, H., Thompson, K., Pierschbacher, M.D., and Ruoslahti, E. (1987) J.Cell Biol. 105, 1183-1190
  - Corbi, A.L., Miller, L.J., O'Connor, K., Larson,
     R.S., and Springer, T.A. (1987) EMBO J. 6, 4023-4028
  - 13. Briesewitz, R., Epstein, M.R., and Marcantonio, E.E. (1993) J.Biol.Chem. 268, 2989-2996
- - 15. Hogervorst, F., Kuikman, I., van Kessel, A.G., and Sonnenberg, A. (1991) Eur.J.Biochem. 199, 425-433
- 30 16. Takada, Y. and Hemler, M.E. (1989) J.Cell Biol. 109, 397-407
  - 17. Takada, Y., Murphy, E., Pil, P., Chen, C., Ginsberg,
     M.H., and Hemler, M.E. (1991) J.Cell Biol. 115,
     257-266
- 35 18. Van der Vieren, M., Le Trong, H., Wood, C.L., Moore, P.F., St.John, T., Staunton, D.E., and Gallatin, W.M. (1995) Immunity. 3, 683-690

- Schnapp, L.M., Breuss, J.M., Ramos, D.M., Sheppard,
   D., and Pytela, R. (1995) J.Cell Sci. 108, 537-544
- 20. Shaw, S.K., Cepek, K.L., Murphy, E.A., Russell, G.J., Brenner, M.B., and Parker, C.M. (1994) J.Biol.Chem. 269, 6016-6025
- 21. Suzuki, S., Argraves, W.S., Arai, H., Languino, L.R., Pierschbacher, M.D., and Ruoslahti, E. (1987) J.Biol.Chem. 262, 14080-14085
- 22. Ignatius, M.J., Large, T.H., Houde, M., Tawil, J.W.,
  Barton, A., Esch, F., Carbonetto, S., and Reichardt,
  L.F. (1990) J.Cell Biol. 111, 709-720
  - 23. Gullberg, D., Gehlsen, K.R., Turner, D.C., Åhlén, K., Zijenah, L.S., Barnes, M.J., and Rubin, K. (1992) EMBO J. 11, 3865-3873
- 15 24. Staaz, W.D., Rajpara, S.M., Wayner, E.A., Carter, W.G., and Santoro, S.A. (1989) J.Cell'Biol. 108, 1917-1924
  - 25. Holmvall, K., Camper, L., Johansson, S., Rubin, K., Kimura, J.H., and Lundgren-Akerlund, E. (1995)
- 20 Exp.Cell Res. 221, 496-503
  - 26. Forsberg, E., Ek, B., Engström, Å., and Johansson, S. (1994) Exp.Cell Res. 213, 183-190
  - 27. Wayner, E.A. and Carter, W.G. (1987) J.Cell Biol.
    105, 1873-1884
- 25 28. Weitzman, J.B., Pasqualini, R., Takada, Y., and Hemler, M.E. (1993) J.Biol.Chem. 268, 8651-8657
  - 29. Elices, M.J. and Hemler, M.E. (1989)
    Proc.Natl.Acad.Sci.U.S.A. 86, 9906-9910
- 30. Languino, L.R., Colella, S., Zanetti, A., Andrieux,
  30 A., Ryckewaert, J.J., Charon, M.H., Marchisio, P.C.,
  Plow, E.F., Ginsberg, M.H., Marguerie, G., and et al
  (1989) Blood 73, 734-742
  - 31. Tuckwell, D.S., Humphries, M.J., and Brass, A. (1994) Cell Adhes.Commun. 2, 385-402
- 35 32. Springer, T.A. (1997) Proc.Natl.Acad.Sci.U.S.A. 94, 65-72

- 33. Colombatti, A., Bonaldo, P., and Doliana, R. (1993)
  Matrix 13, 297-306
- 34. Lee, C.H., Bradley, G., and Ling, V. (1995) Cell Growth Differ. 6, 347-354
- 5 35. Calderwood, D.A., Tuckwell, D.S., and Humphries, M.J. (1995) Biochem.Soc.Trans. 23, 504S
  - 36. Kern, A., Eble, J., Golbik, R., and Kuhn, K. (1993) Eur.J.Biochem. 215, 151-159
- 37. Tuckwell, D.S., Reid, K.B., Barnes, M.J., and
  Humphries, M.J. (1996) Eur.J.Biochem. 241, 732-739
  - 38. Kamata, T. and Takada, Y. (1994) J.Biol.Chem. 269, 26006-26010
  - 39. Dürr, J., Goodman, S., Potocnik, A., von der Mark, H., and von der Mark, K. (1993) Exp.Cell Res. 207, 235-244
  - 40. Salter, D.M., Hughes, D.E., Simpson, R., and Gardner, D.L. (1992) Br.J.Rheumatol. 31, 231-234
- 41. Woods, V.L.J., Schreck, P.J., Gesink, D.S., Pacheco, H.O., Amiel, D., Akeson, W.H., and Lotz, M. (1994)

  Arthritis Rheum. 37, 537-544
  - 42. Camper, L., Heinegård, D., and Lundgren-Åkerlund, E. (1997) J.Cell Biol 138, 1159-1167
  - 43. Hemler, M.E., Sanchez Madrid, F., Flotte, T.J.,
    Krensky, A.M., Burakoff, S.J., Bhan, A.K., Springer,
    T.A., and Strominger, J.L. (1984) J.Immunol, 132.
- 25 T.A., and Strominger, J.L. (1984) J.Immunol. 132, 3011-3018
  - 44. Bottger, B.A., Hedin, U., Johansson, S., and Thyberg, J. (1989) Differentiation. 41, 158-167
- 45. Sommarin, Y. and Heinegard, D. (1983) Biochem. J. 214, 30 777-784
  - 46. Häuselmann, H.J., Aydelotte, M.B., Schumacher, B.L., Kuettner, K.E., Gitelis, S.H., and Thonar, E.J.M.A. (1992) Matrix 12, 116-129
  - 47. Miller, E.J. (1972) Biochemistry 11, 4903-4909
- 35 48. Wessel, D. and Flugge, U.I. (1984) Anal.Biochem. 138, 141-143

- 49. Blobel, G. and Dobberstein, B. (1975) J.Cell Biol. 67, 835-851
- 50. Hellman, U. (1997) in Protein structure analysis.

  Preparation, characterization, and microsequencing

  (Kamp, R.M., Choli-Papadopoulou, T., and WittmannLiebold, B., eds) pp. 97-104, Spriner-Verlag,

  Heidelberg
  - 51. Charles, I.G., Palmer, R.M., Hickery, M.S., Bayliss, M.T., Chubb, A.P., Hall, V.S., Moss, D.W., and
- 10 Moncada, S. (1993) Proc.Natl.Acad.Sci.U.S.A. 90, 11419-11423
  - 52. Tuckwell, D.S., Brass, A., and Humphries, M.J. (1992) Biochem.J. 285, 325-331
- 53. Dedhar, S. and Hannigan, G.E. (1996) Curr.Opin.Cell 15 Biol. 8, 657-669
  - 54. Hughes, P.E., O'Toole, T.E., Ylanne, J., Shattil, S.J., and Ginsberg, M.H. (1995) J.Biol.Chem. 270, 12411-12417
- 55. Puzon McLaughlin, W., Yednock, T.A., and Takada, Y. (1996) J.Biol.Chem. 271, 16580-16585
  - 56. O'Toole, T.E., Katagiri, Y., Faull, R.J., Peter, K., Tamura, R., Quaranta, V., Loftus, J.C., Shattil, S.J., and Ginsberg, M.H. (1994) J.Cell Biol. 124, 1047-1059
- 25 57. De Melker, A.A., Kramer, D., Kuikman, I., and Sonnenberg, A. (1997) Biochem J 529-537
  - 58. Rojiani, M.V., Finlay, B.B., Gray, V., and Dedhar, S. (1991) Biochemistry 30, 9859-9866
- 59. Coppolino, M.G., Woodside, M.J., Demaurex, N.,
  30 Grinstein, S., St Arnaud, R., and Dedhar, S. (1997)
  Nature 386, 843-847
  - 60. Hynes, R.O. (1992) Curr.Opin.Genet.Dev. 2, 621-624
  - 61. Santoro, S.A. (1986) Cell 46, 913-920
- 62. Languino, L.R., Gehlsen, K.R., Wayner, E., Carter, W.G., Engvall, E., and Ruoslahti, E. (1989) J.Cell Biol. 109, 2455-2462

WO 99/51639 PCT/SE99/00544

35

63. Yokosaki, Y., Monis, H., Chen, J., and Sheppard, D. (1996) J.Biol.Chem. 271, 24144-24150

SE	QUENC	JE	LTS.	ΙΤŃ	G.																. `
(1	) GE (i)									2	v										
(2	(ii) (ii) (vi)	SI A) 3) C) O) M OI E)	EQUE LEI TYI STI TOI OLEG RIGI ORG	ENCI NGT PE: RAN POL CUL INAI GAN LLT	E C H: DEI OGY AR L S YPE	HAI 38 acl ONE C: TY OU!	RAC 84 ei SS li PE RCF hui	b c : () me:	ERI ase aci dou ar cDi	ST pld ld lbl	IC: an e	s: rs d	am:		-						
٠.	1		GTCA(	+-			+				+			-+-			+			+	60
a								M	E	L	P	F	v	T	H ·	L	F	L	P	L	-
	61		TTCC	+-			+-				+			-+-			+			+	120
a		V	F L	T	G	L	С	S	. <b>P</b>	F	N	L	D	E	Н	H	P	R	L	F	
	121		GGGC(	+-			+-				+			-+-			+			+	180
a	,	P	G P	P	E	A	E	F	G	Y	s	v	L	Q	н	<b>v</b> .	G	G	G	Q	-
	181		TGGA'	+-	<b></b>		+		<u>-</u>		+			-+-			+			+	240
a a		R	W M	L	V	G	A	P	W	D	G	P	s	Ġ	D	R	R	G	D	V	<u>.</u> .
			CGCT																		
	241		GCGA																		300
a		Y	R C	P	٧	G	G	A	H	N	A	P	С	A	K	G	Н	L	G	D	-
	. 301		CAAC																		360
			COTO C																		

Y Q L G N S S H P A V N M H L G M S L L  ${\tt GAGACAGATGGTGATGGGGGATTCATGGCCTGTGCCCCTCTCTGGTCTCGTGCTTGTGGC}$ CTCTGTCTACCACTACCCCCTAAGTACCGGACACGGGGAGAGACCAGAGCACGAACACCG E T D G D G G F M A C A P L W S R A C G

	401		CTC	TGT	CTT	CAG	TTC	TGG	GAT.									CCA			GGGA	480
	421		GAG	ACA	GAA	GTC	AAG	ACC	CTA												CCCT	400
a		s	s	v	F	s	s	G	I	С	A	R	V	D	A	S.	F	Q	P	Q	G	-
	481				-+-			+				+			-+-			+			GGAT	540
		TC	GGA	CCG	TGG	GTG	ACG	GGT	TGC	GAC	GGG	TTG	TAT	GTA	CCT	ACA.	ACA	GTA	ACA	GAA	CCTA	
a		S	L	A	P	T	A	Q	R	С	P	T	Y	M	D	V	V	I	V	L	D	-
	541				-+-			+			<b>-</b>	+			-+-			+				600
									٠												rccc	
a ·		G	S	N	S	I	Y	P	W	S	Е	V	Q	Т	F	Ţ.	R ·	R	L	V	G	-
	601																				CCCT	660
		TT	TGA	CAA	ATA	ACT	GGG	TCT	TGT	CTA	TGT	CCA	CCC:	TGA	CCA	rgr	CAT	ACC	CCT	CTC	GGGA	
a		ĸ	L	F	Ţ	D	P	Ē	Q	I	Q	V	G	L	V	Q	. <b>Y</b>	G	E,	s	P	-
	661				-+-			+				+~-			-+-			+				720
																_					TTTC	
a		V	H	E	W	S.	L	G.	D ·	F	R	T	K	E	Ε.	V	V	R	A	A	K	-
	721				-+-			+				+			-+-			+			CTGC + SACG	780
a´		N	L	s	R	R	E	G	R	E	T	K	T	A	Q	A	I	M	V.	A	С	-
		AC	AGA	AGG	GTT	CAG	TCA	GTC	CCA	TGG	GGG	CCG.	ACC	CGA	GGC:	rgce	CAG	GCT2	ACT	GGT	GGTT	
	781													•			-		rga	CCA	+ CCAA	840
a		T	E	G	F	s	Q	S	Н	G	G	R	P	E	A	A	R	L.	L	v	v	-
	841																				rgag	
		CA	GTG.	ACT.	ACC	TCT	CAG	GGT	ACT.	ACC:	TCT	CCT	CGA	AGG	ACG:	rcg'	rga:	rtt(	CCG	GAC	ACTC	•
a		<b>V</b>	T	D	G	E	S	H	D	G	E	E	L	P	A	A	L	Ķ	A	C,	E	-
•	901																				GCGA	960
		CG	ACC	TTC	TCA	CTG	TGC	GAT	ACC	CTA	ACG'	TCA	GGA	ACC	AGT	GAT	GGA(	GGC	CGC	CGT	CGCT	
a		A	G	R	V	T	R	Y	G	I	A	V	L	G	H	Y	L	R .	R	Q	R ·	-
	961																				ATTC	1020
																					FAAG	
a		D	P	s	s	F	L	R.	E <sub>.</sub>	I	R	T	I	A	s	D	P	D	E	R	F	-
	1001																				rcgg	1080
	1021																				AGCC	1080
a		F	F	N	v.	T	D	E	A	A	L	Т	D	I	V.	ם	A	L	G	D	R	-

	1081															CTT	TGG	GCT			GTCT	1140
,													GCT			gaa	ACC	CGA			CAGA	1140
a		I	F	G	L	E	G	s	H	A	E	N	E	s	s	F	G	L	E	М	s.	-
																					GGCC	
	1141																				CCGG	1200
a		Q	ı	Ġ	F	s	T	Н	R	L	ĸ	D	G	1	L	F	G	M	v	G	A	-
																					ACGA	
	1201																				+ TGCT	1260
a		Y	D	W	G	G	s	V	L	W	L	E	G	G	Ĥ	R	Ľ,	F	P	P	R	_
																					TTAC	
	1261																				AATG	1320
a		M	A	L	E	D	E	F	P	P.	A	L	Q	N	н	A	A	Y	L	G	Y	_
																					TCGA	
	1321																,				AGCT	1380
a		s	v	s	S	M	L	L	R	G	G.	R	R	L	F	L	s	G	A	P	R	<u> </u>
																					GGŤT	
	1381												CGA		-						CCAA	1440
a		F	R	н	R	G	ĸ	v	·I	A	F	Q	L	ĸ	ĸ	D	G	A	v	R	V	-
																					ATTG	
	1441																				+ FAAC	1500
a		A	Q	s	L	Q	G	E	Q	I	G	s	Y	F	G .	s	Ē	L	С	P	L	-
					TAG	GGA	TGG	AAC													GGA	
	1501				-+- ATC	CCT.	ACC	+ TTG													CCCT	1560
a		D	T	D	R	D	G	T	<b>T</b> -	D	V,	L	L	v	A	A	P	M	F	L	G	_
	15.50	CC	CCA	GAA	CAA	gga.	AAC	AGG													GCTG	
	1561	GG	GGT	CTT	-+- GTT	CCT	TTG	+ TCC													CGAC	1620
a		P	Q	N	ĸ	E	T.	G.	R	v	Y	v	Y	L	v	G	Q	Q	s	L	L	-
	1601																				CATG	
	1621																				TAC	1680
a		T	L	Q	G	т	L	Q	P	E	P	P	Q	D	A	R.	F	G	F	A	М	_
٠	160:																				CTG	
	1681																				AGAC	1740
2		G	Δ.	7	Б		<b>,</b> .				_	_			17	7		~		_		

	1741	GA	AGA'	TGG	GCA(	CCA	GGG	AGC.	ACT(	GTA	CCT	STA(	CCA'	TGG	AAC	CCA	GAG'	TGG	AGTO	CAG	CCC	1800
	1741	CT	TCT	ACC	CGT	GGT	CCC'	TCG	TGA	CAT	GGA	CAT	GGT	ACC	rtg	GGT	CTC.	ACC'	CAC	STC	CGGG	1000
a		E	D	G	H	Q	G	A	L	Y	L	Y	H	Ģ	T	Q	s	G	V	R	P	_
	1801		rcć:	TGC	CCA	GAG	GAT		rgc:							CCŤ	CAG	CTA	CTT1	rggo	CCGA	1860
			AGG	ACG	GGT	CTC	CTA									GGA	GTC	GAT	AA	ACCO	GCT	
a		Н	P	A	Q	R	Ī.	A	A	A	s	M	Ė	Н	A	L	s	Y	ŗ	G	R	-
	1861																	GGC			rgcc	1920
																					ACGG	2,20
a		S	<b>V</b> .	D	G	R	L	D	L	D,	G	D	D .	L	v	D	V	A	V	G	A	<b>-</b> ,
	1921	CA	GGG	GGÇ	AGC	CAT	CCT	GCT	CAG	CTC	CCG		CAT'	_				CCC	ATC	ACTO	GAG	1980
	-3	GT	ccc	CCG'	rcg	GTA	GGA	CGA	GTC(	GAG	GGC	-						GGG:	ragi	rgac	CCTC	
a		Q	Ġ	A	A	I	L	L	S	s	R	P	I	v	Н	L	Т	P	S	L	Ε	-
	1981																				AGCA	2040
																					CGT	
a		V	T .	P	Q	A	Ī	s	V	v	Q	R	Ď	С	R.	R	R	G ·	Q	E	A	-
	2041		CTG	TCT(	GAC'	rgc.	AGC		TTG						CCG	rac'	TCC	TGG:	CGC	CTG	GAT	2100
		CA	GAC	AGA	CTG	ACG'	TCG	GGA	AAC	GAA(	GGT:	rca(	CTG	GAG	GGC.	ATG.	AGG	ACC	AGCC	GAC	CCTA	
a	÷	V	С	L	T	A	A	L	С	F	Q	V	T	s	R.	T	P	G	R	M	D	-
	2101																				rgca +	2160
		GT	GGT'	TAA	GAT(	GTA	CTC	CAA	GTG(	GCG'	rag:	rga	CCT	ACT'	TAC	CTG.	ACG.	ACC	CCG1	rgcz	ACGŢ	
a		H	Q	F	Y	М	R	F	T	A	S	L	D	E	W	T	A	G	A	R	A	
	2161				-+-			+				+			-+-			+-				2220
		CG	TAAI	ACT	ACC	GAG	ACC	GGT	CTC(	CAAC	CAG	GGG2	AGC	CTC	CGA	GGC	CGA	GTC	ACAC	ccc	CTTA	
a		A	F	Đ	G		G .	Q	R	L.	s	P	R	R	L	R	L	Ş	V	G	N	-
	2221				-+-			+				+			-+-			+-				2280
																					CAC	
а				*						•											V	<del>-</del> .
	2281				-+-			+	<u>-</u>			+			-+-			+-				2340
							•							•							ACTC	
a																					E	-
	2341				-+-			+				+			-+-			+-				2400
3							•														N N	_
a		G	3	Ľ	L	۵		v	V	L	v	Ľ	Ľ	J	L	IJ	L	G	2	υ	[A	-

	2401																				GGCC	2460
	2401																				CCGG	2400
a		E	С	V	T .	D	L	v	L	Q	v	N	M	D	I	R	G	s	R	ĸ	A	-
	2461																				CAGA	2520
	2401																				GTCT	2320
a .	-	P	F	v	v	R	G	G	R	·R	ĸ	v	L	v	s	T	T	L	E	N	R	
	2521			•																	GGCC	2580
	2321																				CCGG	2580
a _		ĸ	E	Ň	A	Y	N	T	s	L.	s	I	I	F	s	R	N	L	н	L	A	-
•	. 2501																				CAT	0510
	2581																				AGTA	2640
a		s	L	T	P	Q	R	E	s	P	I	ĸ	v	E	C	A	A	P	s	Ά	н	-
	2641																				rctg	
	2641																				AGAC	2700
a		A	R	L	c	s	V	G	H	P	v	F	Ω.	T	G	A	ĸ	v	T	F	L	
																					rgcc	
	2701																				ACGG	2760
a		L	E	F	E	F	s	С	s	s	L	L	s	Q	v	F	G	ĸ	L	T	A	-
	07.61																				AGCC	
	2761																				rcgg	2820
a		s	s	D	s	L	E	R	N	G	T	L	Q	E	N	T	A	Q	T	s	A	-
	2001																				rgag	
	2821																				CTC	2880
a		Y	I	Q	Y	E	P	H	L	L	F	s	s	E	s	T	L	H	R	Y	Ε.	-
	2001																				CAGG	. ,
	2881						•														TCC	2940
a ´		٧.	н	P	Y	G	T	L	P.	v	G	P	G	P	E	F	ĸ.	T	T	Ľ.	R	-
																					AGCT	
	2941																		-		CGA	.3000
a		v	Q	N	L	G	С	Y	v	v	s	G	Ľ	Ĭ	I	s	A	L	L	P	Α .	- ,
	2021	GT	GGC	CCA	TGG	GGG	CAAT	rta(	CTT	CCT	ATC	ACTO	STCT	CAZ	AGT	CAT(	CAC'	TAA	CAAT	GCF	AAGC	
	3001																				TCG	3060
a		v	A	Н	Ģ	G	N	Y	F	L	ś	L	s	Q	v	I	Т	N	N	А	S	

	3061											AGG +									TCAA +	3120
																					AGTT	3120
a .		С	I	V	Q	N	L.	T	E	P	P.	G	P	P	V	Н	P	E	E	. L	Q	-
		CA	CAC	AAA	CAG	ACT	GAA	TGG	GAG	CAA	TAC	TCA	GTG'	rca	GGT	GGT	GAG	GTG	CCA	CCT	TGGG	
	3121																				+ ACCC	3180
_																						
a		н	T	N	Ķ		N	G	s	Ŋ	Т	Ų.	L	Q	٧	٧		C	п	ы	G	-
	3181																				ATTT +	3240
	•																				TAAA	
a		Q	Ĺ	A	K	G	T	E	ŢV	s	V	G	L	L	R	L	V	Н	N	E	F	-
	2041																				CGAA	2200
																					GCTT	3300
a		F	R	R	A	ĸ	F	ĸ	s.	L	T	v	v	s	T	F	E	L	G	т	E .	-
	-	GA	GGG	CAG	TGT	CCT	ACA	GCT	GAC	TGA	AGC	CTC	CCG'	ГТG	GAG	TGA	GAG	ССТ	СТТ	GGA	GGTG	
	3301																				+ CÇAC	3360
_			•					٠.													V	
а							_		T				٠.				s			_		-
	3361																				AGGG +	3420
	-	CA	AGT	CTG	GGC	CGG	ATA	GGA	GTA	GAG	GGA	CAC	CTA	GGA(	GTA	TCC	GTC.	ACA	GGA	CCĊ.	TCCC	
a		V	Q	T	R	P	I	L	I	S	L	W	I	L	Ľ	G	S	V	L	G <sub>.</sub>	G	-
	3421																				TAAG	3480
																					ATTC	3400
a ·		L	L	· L	ŗL	A	L	L	v	F	C:	L	W	ĸ	Ļ	G	F	F	A	H	ĸ	- , '
																					GGGT	
	3481																				CCCA	3540
a		ĸ	I	P	E	E	E	ĸ	R	E	Е	ĸ	L	E		-						
		CT:	AGA	DAG	TCC	TCC	стс	CCA	GCT'	ጥጥረት	יייירי	שמכ	מממ	اششت	CCA	ממד	AAG	CAG	AGG	ጥጥጥ.	GGGG	-
	3541				-+-			+				+			-+-			+			+	3600
•		GA	rcr	TTC	AGG	AGG	GAC	CGT	CGA	AAG	AAG	TTC	rcr	зАА	CGT	ATT	rrc	GŢC	TCC	AAA	CCCC	
•																					TTGA	
	3601																				AACT	3660
	3661																				GAGC	3720
	2001											•									CTCG	3720

3721	TGGCACCAAAACTAGCCATGCTCCCACCCTCTGCTTCCCTCCTCGTGATCCTGGTTC	2200
3/21	ACCGTGGTTTTGATCGGTACGAGGGTGGGAGACGAAGGGAGGAGGAGCACTAGGACCAAG	3780
3781	CATAGCCAACACTGGGGCTTTTGTTTGGGGTUCTTTTATCCCCAGGAATCAATAATTTTT	3840
3841	TTGCCTAGGAAAAAAAAGCGGCCGCGAATTCGATATCAAGCT	

(2)	IN	FO	RMA	ΙΤ	ON	F	OR	SE	Q :	ΙD	NC	).	2:						•			
	( :	Ĺ)	S	EQ	UE	NCE	G C	HΑ	RA	CTE	ERI	ST	IC	S:								
				(A)	]	LEN	IGT	Н:	3	779	b	as	e p	pai	rs							
				(B)		ΓYΕ	E:	n	uc]	lei	.d	ac	id	an	d	am.	ino	a	ci	d		
				(C)		STF	NAS	DE	DNE	ess	::	do	ub]	lė				•				
				(D)		roe	OL	OG	Υ:	li	.ne	ar										
				(E)	ı																	
	( :	L)		IOL		ULZ	λR	ΤY	PΕ	: 0	:DN	ΙA			-							
		-		RI																		
	.``			(A)							ıım:	an					•					
												ndı		·v+	_							
				(2)						Ŭ				., .	_							
	. (	zi'	۱ ۶	EO	HE:	NCF	. г	ES	ĊR:	רק ד	ידכ	N:	S	EO:	ID	N	ά.	2.		•		
	( 2	<b>.</b> ,	, -	υLQ	01.			,	O1(.					υQ	10	. 11	٠.		ļi			
•																						
																					CTG	
	1																				+	
	."	GT	CCA	GTC	T.L.	GGC	IAG	rec	3 I A	CCT	IGA	افافافا	GAA	GCA	3 I G/	1GT (	3GA(	AA	JAA(	اجاحات	GAC	
						:		*	М	E	L	P	F	V	T	H	L	F	L	P	L	-
		~=							~~~	~~~			~ ~ m	~~~	<b>70.5</b>					- com		
	61																				ATTC +	120
																					raag	
					_			_		_	_		_		_			_	_			
a		V	F	L	T	G	L	С	S	Р	F.	N	ь	D	Ε	Н	н	Þ	R	ь	F.	-
		CC	AGG	GCC	ACC.	AGA.	AGC'	TGA	ATT'	TGG	ATA	CAG	TGT	CTT	ACA	ACA:	rgt:	rgg	GGT	rggz	ACAG	
	121																				+	180
		GG	TCC	CGG'	TGG	TCT"	TCG.	ACT:	PAA	ACC'	rat	GTC.	ACA	GAA'.	rGT'	'GT'	ACA	4CC(	CCC	ACC"	rgtc	
a		P	G	P	P	E	Α	E	F	. G	Y	s	V	L	Q	H	v ·	G	G	G	Q	-
										, 												
	181																				CGTT	240
	101																				CAA	
		_					_	_	_		_	_	_	_	_			_	_	_	••	
a		R	W	M	ь	٧	G	A	Р	W,	D	G	Р	S	G	D	K	R	G	D	V	
		TA	TCG	CTG	CCC	TGT.	AGG	GGG	GGC	CCA	CAA	TGC	ccc	ATG:	rgc	CAAC	GGG	CA	CTT	AGG!	rgac	
	241																					300
		AT	AGC	GAC	GGG	ACA	TCC	CCC	CCG	GGT	GTT.	ACG	GGG'	TAC	ACG	STTC	CCC	3GT(	SAA'.	rcc.	ACTG	
a		Y	R	C	P	<b>v</b> .	G	G.	A	н	N	A	P	С	A	K	G	Н	L	G	D	_
															<u>.</u>							
	301																				ATTE	360
																					CAAT	300
a		Y	Ò	L	G <sub>.</sub>	N	S	s	H	P	A	V	N	M	H	L	G	M	s	L.	L	-
		GA	GAC	AGA'	TGG	TGA	TGG	GGG.	ATT	CAT	GGC	CTG	TGC	CCC'	rcro	CTG	GTC:	r¢g:	rgc:	rtg	rggc	
	361																				+	420
		CT	CTG	TCT.	ACC.	ACT	ACC	CCC'	TAA	GTA	CCG	GAC	ACG	GGG	AGA	SAC	CAG	AGC	ACG	AAC	ACCG	
a		E	Т	D	G	D	G	G	F	М	A	С	A	P	L	W	s	R	A	Ċ	G .	<del>-</del>
4											•								•			-
	421																				GGA	480
																					CCT	

a		s s	s v	F	S	S	G	I	С	A	R	V	D	A	S	F	Q	P	Q	G	-
	481		CTGG	CACC	CAC	TGC	CCA					ATA						TGT	CTT	GGAT	540
	101		GACCO	STGG	GTG	ACG	GGT											ACA	GAA	CCTA	
a		s i	. A	P	T	A	Q	R	С	P	T	Y	M	D	V.	V	I	V	L	D <sub>.</sub>	-
	E 41		CCA																	AGGG	600
	341		AGGTT																		
а		G	s N	s	1	Y	P	W	s	E	v	Q	T	F	L	R	R	L	v	G	<u>-</u>
			CTGTT	rat'i	'TGA	ccc	AGA										TGG	GGA	GAG		
	601		ACA!	ATA	ACT	'GGG	TCT					CCC					ACC	CCT	CTC		660
a		ĸ 1	F	I	D	P	E	Q	I	Q	v	G	L	v	Q	Y	G	E	s	P.	-
			CATG																		
	661	CATO	TAC																		720
a		V I	i E	w	s	L	G	D	F	R	T	ĸ	E	E	v	v ·	R	A	A	ĸ	-
			CTCAC	STCG	GCG	GGA	.GGG												GGC	CTGC	-
	721		SAGTO	CAGO	CGC	CCT	CCC				•	CTG							CCG	GACG	780
а		N I	s s	R	Ŕ	E	· G	R	E	T	K	T	A	Q	A	I	м	v	A	С	-
			SAAGO																	GGTT	
	781	TGT	CTTC																	CCAA	840
a		T I	G G	F	s	Q	s	Н	G	G	R	P	E	A	A	R	L	L	v	v	-
	0.41		ACTG															GGC			000
	. 841		GACI															CCG		ACTC	900
a		v :	r D	Ğ	E	s	н	D	. <b>G</b>	.E	E	L	P	A	A	L	ĸ	A	c	E.	<b>-</b> .
																	CCG	GCG	GCA	GCGA	
	901	CGA	CTT											-			GGC	CGC	CGT		960
a		A (	3 R	v	T	R	Y	G	I	A	v	r	G	н	Y	L	R	R	Q	R	-
			CCAC																		
-	961		GGT																		1020
a		D I	? \$	s	F	L	R	E	I	R	T	I	A	s	D	P	Ď	E	R	F	-
			TCAZ																		٠
		AAG																			1080
a		F 1	e n	v	T	D	E	A	A	L	T	D	ı	v	D	A	L	G	D	R <sub>.</sub>	_
			TTGC																		
	1081																			+ CAGA	1140

a		1.	F	G	L	E	G	S	Н	A	E	N	E	S	S	F	Ģ	L	E	M	S	- 1
	1141		;		-+-			<del>+</del> -			;	<b></b>			+			+				1200
		GT	CTA	ACC	AAA	GAG	GTG	AGT	AGC	CGAT	PŢT(	CCTA	ACCO	TA	\GA/	AA.	4CC	CTA	CCA	CCC	CCGG	
а		Q	I	G	F	s	T	н	R	L	K	D	G	I	L	F	G <sub>.</sub>	M	<b>V</b> .	G	A	-
																					ACGA	
	1201																				rgct	1260
a		Y	ď.	W .	G	G	s	V	L	W	L	E	G	Ġ.	Н	R	L	F	P	Ρ.	R	-
	1261																				TAC	1320
																					ATG	1020
a		М	A	L	E.	D	E	F	P	P	A	L	Q	N	H	A	A	Y	L	G	Y	-
	1 201																				CGA	
	1321																				GCT	1380
a		s	<b>v</b> .	s	s	M	L	L.	R	G	G	Ŗ	R.	L	F	L	s	G	Ä	P	R	-
	1 2 0 1																				GTT	
	1381																				CAA	1440
ạ	,	F	R	H	R.	G	K	v	ı.	A	F.	Q	L	ĸ	K	D.	G	A	v	Ř	V	<b>-</b> .
	1441																				TTG	1500
	7441																				AAC	1300
a		A	Q	s	L	Q	G .	E	Q	I	G	s	Y	F	G	s	E	L	С	P	L	-
	1501																				GGA	1560
	1301			-																	CCT	1360
a		D	T	D	R	D	G	T	Т	D -	v	L	L	V	A	A	P	M	F	L	G .	-
																					CTG	
٠	1561																				GAC	1620
а		P.	.Q	N	ĸ	E	T	G	R	v .	Y	v	Y	L	v	G	Q	Q	s	L	L	_
	1601																				ATG	1.500
	1621																				TAC	1 <b>.</b> 80
a		T	L,	Q.	G	T	L	Q	P	E	P	P	Q	D	A	R	F	G.	F	A	М	-
	1681																				CTG	1740
	100I																				ĠAC	1,40
a	ė	G	A	L .	P.	D	L	N	Q	D	G	F	A	D	V	A	۷.	G	A	P	L	<del>-</del>
	1741																				CCC	1000
	- 11																				rece	1000

а		E	D	G	H	Q	G	A	Ĺ	Y	L	Y	Н	G	T	Q	S.	G	V	R	P	-
	1001																				CCGA	1000
	1801																				GGCT	1860
a		H	P	A	Q-	R	I	A	A	A	s	M	P	H.	A	L	s	Y	F	G	R	-
																					TGCC	
	1991																				ACGG	1920
a		s	v	D	G.	R	L	D	L	D	G	D	D·	L	v	D	v	A	v	G	A	
	1 921																				GGAG	1000
	1921																				CCTC	1980
а		Q	G.	A	A	I	L	L	s	s	R	P	I	v	H	L	T	P	s	L	Ē	-
	1981																				AGCA	
	1901														-						TCGT	2040
a		v	. <b>T</b>	P	Q	A	I	s	V	v	Q	R	Đ	C	R	R	R	Ġ	Q	E.	A	-
	.2041																				GGAT	2100
						,															CCTA	2100
a		V	С	L	<b>T</b> .	A	A	L	С	F	Q	<b>v</b> .	T	8	R	T	P	G	R	W	D.	-
	2101																				rgca	2160
		GT	GGT	'TAA	.GAT	GTA	CTC	CAA	GTG	GCG	TAG	TGÁ	CCT	ACT	TAC	CTG	ACG	ACC	CCG	IGC.	ACGT	
a		H	· Q	F	Y	M .	R	F	T	A	S	L	D	E	W	T	A	G	A	R	Α	-
	2161			TGA																	SAAT	2220
		CG	TAA	ACT	ACC	GAG	ACC	GGT	CTC	CAA	CAG	GGG	AGC	CTC	CGA	GGC	CGA	GTC.	ACA	CCC	CTTA	
а		A			G		-	Q		L							L	S	V	G	N	-
	2221				-+-			+				+			-+-			+				2280
																				CGG'	rcac	
a			_	-	. E	-			_		÷	_	-	T	s		Y	L	R	P	V	<b>-</b> . ·
	2281				-+-			+			- <del>-</del> -	+			-+-			+				2340
-															•						ACTC	
a		:									• •										E	
	2341				-+-			+				+			-+-			+				2400
a		G	•		Т																STTA	_
a		-					•	_		•						•		G CTIC				-
	2401				-+-			+				+			-+ <b>-</b>			+			GGCC + CCGG	2460
		- L	100	$\alpha$	U112	101	ᄱᄱ	-	-CA	L D C	ILM	-11	MIN.		יאניט		ょしい	أحالتك	316	-17		

а	•	E	С	V 1	r D	L	v	L	Q	V	N	M	D	I	R	G	S	R	K	A	-
	0461				GTTC																2520
	2461				CAAG																2520
a	-	P	F	v v	V R	G	G	R	R	K	v	L	v	s	T	T	L	E	N	R	
					GCTT																
	2321				CGAA'																2580
а		ĸ	Ē	N Z	A Y	N	T	s	L.	s	I	I	F	s	R	N	L	Н	L	A	-
	2501				CCTC																2640
	2301	-			GGAG'																2040
а		S	F	T I	P Q	R	E	s	P	I	ĸ	V	E	,c	A	A.	₽.	s	A	H.,	<b>-</b> ,
	2641				TGCA																2700
	2041				ACGT																2700
a ·	•	A	R	L (	c s	V	G	Н	P	v	F	Q	T	G	A	ĸ	V.	T	F	L	
	2701				GAGT1																2760
-	2.01				ETCA					-											2,00
a		L	E	F I	E F	s	С	s	S	L,	L	S	Q	V	F	G	K	L	T	A	<del>-</del> .
	2761				AGCCI																2820
					rcgg/																
a	• .	S	S	D S	S L	·E	R	N	G	T	L	Q	E	N	T·	A	Q	T .	s	A	-
	2821				FATG																2880
					ATACI																
a .		Y	I	Q Y	Y E	P	Н	L	L	F	s	s	E	s	T	L	Н	R	Υ.	E	-
	2881				ratgo +																2940
		CA	AGTG	GGT	ATAC	CCTG	GGA	GGG	TCA	ccc	AGG	ACC	GGG	TCT'	TAA	GTT:	rtg	GTG/	AGA(	STCC	
a		V	H	P Y	Y G	T	L	P	V	G	P	G	P	E	F	K	T	T	L	R. ·	-
	2941				GCAA(																3000
					CGTT																,
а		T	N	N A	A S	С	I	V	Q	N	L	T	E	P	P	G	P	P	V	Н	-
	3001				CTTC#																3060
					GAAG!																
a		P	E	E I	L Q	H	T	N	R	L	N	G	s	N	T	Q.	C	Q	V	V	-
	3061				CTTG(																3120
																				CGAC	2120

a		R.	C	H	L	G	Q	L	A	K	G	T	E	V	S	V	G	L	L	R	L	-
		GTTCACAATGAATTTTTCCGAAGAGCCAAGTTCAAGTCCCTGACGGTGGTCAGCACCTTT													3180							
a			AGT H				•				STT(					T T		CCA V	GTC S	GTG( T	GAAA F	_
-					-+-			+			+	<b></b> -			-+			+			TGAG + ACTC	3240.
a				·G																	Ė	<b>.</b> .
	3241				-+-			+		;	4	<b></b> -			-+			+			AGGC + ICCG	3300
a		s		L		v					P		L,		s	L	W	I	L	I	G	'-
	3301				-+-			+		·		<b></b> -			-+			+			TGGC + ACCG	
a				L.																	G	<b>-</b> ,
	3361				-+-			+			+	<b></b> -			+		1_	+			ATGA + FACT	3420
а		F	F	A	H	K	K	I	P	E	E	Ë.	K	R	E	E	K	L	E	Q		
	3421				-+-			+		<b>-</b> -	+	<b>+</b> -			-+			+			TAAA + ATTT	3480
	3481				-+-			+				+			+			+			GACC + CTGG	3540
	3541				-+-			+				+			-+-			+			TACC + ATGG	3600
	3601				-+-			+		-,		+			-+			+			CTCC + GAGG	3660
	3661				-+-			+				+			-+-			+			CCAG + GGTC	3720
	3721			AAT.	-+-			+				+			-+			+			;	3779

(2)	INFORMATION	FOR	SEQ	ID	NO.	3:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 143 base pairs
  - (B) TYPE: nucleic acid and amino acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (iii) MOLECULAR TYPE: cDNA
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: human
  - (B) CELLTYPE: chondrocyte
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 3:

NdeI

GGGGCATATGGTTCAGAACCTGGGTTGCTACGTTGTTTCCGGTCTGATCATCTCCGCTCT

1 -----+ 60
CCCCGTATACCAAGTCTTGGACCCAACGATGCAACAAAGGCCAGACTAGTAGAGGCGAGA

b - G H M V Q N L G C Y V V S G L I I S A L -

GCTGCCGGCTGTTGCTCACGGTGGTAACTACTTCCTAAGCTTGTCCCAGGTTATCAGCGG
61 -----+ 120
CGACGGCCGACAACGAGTGCCACCATTGATGAAGGATTCGAACAGGGTCCAATAGTCGCC

b LPAVAHGGNYFLSLSQVISG-

BamHI

b LVPRGSP -

25

## CLAIMS

- 1. A recombinant or isolated integrin subunit  $\alpha 10$  comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or homologues or fragments thereof having similar biological activity.
- 2. A process of producing a recombinant integrin subunit  $\alpha$ 10 comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or homologues or fragments thereof having similar biological activity, which process comprises the steps of
- a) isolating a polynucleotide comprising a nucleotide sequence coding for an integrin subunit  $\alpha 10$ , or homologues or fragments thereof having similar biological activity,
  - b) constructing an expression vector comprising the isolated polynucleotide,
- c) transforming a host cell with said expressionvector,
  - d) culturing said transformed host cell in a culture medium under conditions suitable for expression of integrin subunit  $\alpha 10$ , or homologues or fragments thereof having similar biological activity, in said transformed host cell, and, optionally,
  - e) isolating the integrin subunit  $\alpha 10$ , or homologues or fragments thereof having similar biological activity, from said transformed host cell or said culture medium.
- 3. A process of providing an integrin subunit α10, 30 or homologues or fragments thereof having similar biological activity, whereby said subunit is isolated from a cell in which it is naturally present.
  - 4. An isolated polynucleotide comprising a nucleotide coding for an integrin subunit  $\alpha 10$ , or for homologues or fragments thereof, which polynucleotide comprises the nucleotide sequence shown in SEQ ID No. 1 or SEQ ID No. 2 or suitable parts thereof.

30

35

- 5. An isolated polynucleotide or oligonucleotide which hybridises to a DNA or RNA encoding an integrin subunit  $\alpha 10$  or homologues or fragments thereof, wherein said polynucleotide or oligonucleotide fails to hybridise to a DNA or RNA encoding an integrin subunit  $\alpha 1$ .
- 6. A vector comprising a polynucleotide or oligonucleotide coding for an integrin subunit  $\alpha 10$ , or homologues or fragments thereof, which polynucleotide or oligonucleotide comprises the nucleotide sequence shown in SEQ ID No. 1 or SEQ ID No. 2 or parts thereof.
- 7. A vector comprising a polynucleotide or oligonucleotide which hybridises to a DNA or RNA encoding an integrin subunit α10 or homologues or fragments thereof,
   15 wherein said polynucleotide or oligonucleotide fails to hybridise to a DNA or RNA encoding an integrin subunit α1.
  - 8. A cell containing the vector as defined in any one of claims 6 and 7.
- 9. A cell generated by the process in claim 2, in which a polynucleotide or oligonucleotide coding for an integrin subunit α10, or homologues or fragments thereof, which polynucleotide or oligonucleotide comprises the nucleotide sequence shown in SEQ ID No. 1 or SEQ ID No. 2 or parts thereof has been stably integrated in the cell genome.
  - 10. Binding entities having the capability of binding specifically to integrin subunit  $\alpha 10$  comprising the amino acid sequence of SEQ ID No. 1 or SEQ ID No. 2, or to homologues or fragments thereof.
  - 11. Binding entities according to claim 10, which are chosen from the group comprising proteins, peptides, carbohydrates, lipids, natural integrin binding ligands, polyclonal and monoclonal antibodies, and fragments thereof.
  - 12. A recombinant or isolated integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , in which the

15

25

30

35

subunit  $\alpha 10$  comprises the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, and homologues and fragments thereof having similar biological activity.

- 13. A recombinant or isolated integrin heterodimer according to claim 12, wherein the subunit  $\beta$  is  $\beta$ 1.
- 14. A process of producing a recombinant integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , in which the subunit  $\alpha 10$  comprises the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, and homologues and fragments thereof, which process comprises the steps of
- a) isolating one polynucleotide comprising a nucleotide sequence coding for a subunit  $\alpha 10$  of an integrin heterodimer and, optionally, another polynucleotide comprising a nucleotide sequence coding for a subunit  $\beta$  of an integrin heterodimer, or polynucleotides or oligonucleotides coding for homologues or fragments thereof having similar biological activity,
- b) constructing an expression vector comprising said isolated polynucleotide coding for said subunit  $\alpha 10$  optionally in combination with an expression vector comprising said isolated nucleotide coding for said subunit  $\beta$ ,
- c) transforming a host cell with said expression vector or vectors,
- d) culturing said transformed host cell in a culture medium under conditions suitable for expression of an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or homologues or fragments thereof having similar biological activity, in said transformed host cell, and, optionally,
- e) isolating the integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or homologues or fragments thereof having similar biological activity, or the  $\alpha 10$  subunit thereof from said transformed host cell or said culture medium.
- 15. A process of providing an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta\text{,}$  or homologues

25

or fragments thereof having similar biological activity, whereby said integrin heterodimer is isolated from a cell in which it is naturally present.

- 16. A cell containing a first vector, said first vector comprising a polynucleotide or oligonucleotide coding for a subunit  $\alpha 10$  of an integrin heterodimer, or for homologues or parts thereof having similar biological activity, which polynucleotide or oligonucleotide comprises the nucleotide sequence shown in SEQ ID No. 1 or SEQ ID No. 2 or parts thereof, and a second vector, said second vector comprising a polynucleotide or oligonucleotide coding for a subunit  $\beta$  of an integrin heterodimer, or for homologues or fragments thereof.
- 17. Binding entities having the capability of binding specifically to the integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or to homologues or fragments thereof, or a subunit  $\alpha 10$  thereof, having similar biological activity.
  - 18. Binding entities according to claim 17, wherein the subunit  $\beta$  is  $\beta$ 1.
    - 19. Binding entities according to claim 17 or 18, which are chosen among the group comprising proteins, peptides, carbohydrates, lipids, natural integrin binding ligands, and fragments thereof.
  - 20. A fragment of the integrin subunit  $\alpha$ 10, which fragment is a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the spliced domain.
- 21. A fragment according to claim 20, which is a 30 peptide comprising the amino acid sequence KLGFFAHKKIPEEEKREEKLEQ.
  - 22. A fragment according to claim 20, which comprises the amino acid sequence from about amino acid No. 952 to about amino acid no. 986 of SEQ ID No. 1.
- 35 23. A fragment according to claim 20, which is a peptide comprising the amino acid sequence from about

25

amino acid No. 140 to about amino acid no. 337 of SEO ID No. 1.

- 24. A method of producing a fragment of the integrin subunit  $\alpha 10$  as defined in any one of claims 20-23, which method comprises a sequential addition of amino acids containing protective groups.
- 25. A polynucleotide or oligonucleotide coding for a fragment of the integrin subunit  $\alpha 10$  as defined in any one of claims 20-23.
- 10 26. Binding entities having the capability of binding specifically to a fragment of the human integrin subunit  $\alpha$ 10 as defined in any one of claims 20-23.
  - 27. Binding entities according to claim 26, which are chosen from the group comprising proteins, peptides, carbohydrates, lipids, natural integrin binding ligands, and fragments thereof.
  - 28. A process of using an integrin subunit  $\alpha 10$  comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or an integrin heterodimer comprising said subunit  $\alpha 10$  and a subunit  $\beta$ , or a homologue or fragment of said integrin or subunit having similar biological activity, as a marker or target molecule of cells or tissues expressing said integrin subunit  $\alpha 10$ , which cells or tissues are of animal including human origin.
  - 29. A process according to claim 28, whereby said fragment is a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the spliced domain.
- 30. A process according to claim 29, whereby said 30 fragment is a peptide comprising the amino acid sequence KLGFFAHKKIPEEEKREEKLEQ.
  - 31. A process according to claim 29, whereby said fragment comprises the amino acid sequence from about amino acid no. 952 to about amino acid no. 986 of SEQ ID No. 1.
  - 32. A process according to claim 29, whereby said fragment comprises the amino acid sequence from about

WO 99/51639 PCT/SE99/00544

amino acid no. 140 to about amino acid no. 337 of SEQ ID No. 1.

5

20

25

3.0

35

- 33. A process according to claim 28, whereby the subunit  $\beta$  is  $\beta 1.$
- 34. A process according to claim 28, whereby said cells are chosen from the group comprising chondrocytes, smooth muscle cells, endothelial cells, osteoblasts and fibroblasts.
- 35. A process according to any one of claims 28-34, which process is used during pathological conditions involving said subunit  $\alpha 10$ .
  - 36. A process according to claim 35, which pathological conditions comprise damage of cartilage.
- 37. A process according to claim 36, which patho15 logical conditions comprise trauma, rheumatoid arthritis
  and osteoarthritis.
  - 38. A process according to any one of claims 28-34, which is a process for detecting the formation of cartilage during embryonal development.
  - 39. A process according to any one of claims 28-34, which is a process for detecting physiological or therapeutic reparation of cartilage.
    - 40. A process according to any one of claims 28-34, which is a process for selection and analysis, or for sorting, isolating or purification of chondrocytes.
    - 41. A process according to any one of claims 28-34, which is a process for detecting regeneration of cartilage or chondrocytes during transplantation of cartilage or chondrocytes.
  - 42. A process according to any one of claims 28-34, which is a process for in vitro studies of differentiation of chondrocytes.
  - 43. A process of using binding entities having the capability of binding specifically to an integrin subunit  $\alpha 10$  comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or an integrin heterodimer comprising said subunit  $\alpha 10$  and a subunit  $\beta$ , or to homo-

20

25

30

. 35

logues or fragments thereof having similar biological activity, as markers or target molecules of cells or tissues expressing said integrin subunit  $\alpha 10$ , which cells or tissues are of animal including human origin.

- 44. A process according to claim 43, whereby said fragment is a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the spliced domain.
- 45. A process according to claim 43, whereby said fragment is a peptide comprising the amino acid sequence KLGFFAHKKIPEEEKREEKLEQ.
  - 46. A process according to claim 43, whereby said fragment comprises the amino acid sequence from about amino acid no. 952 to about amino acid no. 986 of SEO ID No. 1.
  - 47. A process according to claim 43, whereby said fragment comprises the amino acid sequence from about amino acid no. 140 to about amino acid No. 337 of SEQ ID No. 1.
  - 48. A process according to claim 43, whereby the subunit  $\beta$  is  $\beta1$ .
  - 49. A process according to any one of claims 43-48, which is a process for detecting the presence of an integrin subunit  $\alpha 10$  comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or of an integrin heterodimer comprising said subunit  $\alpha 10$  and a subunit  $\beta$ , or of homologues or fragments thereof having similar biological activity.
  - 50. A process according to any one of claims 43-48, which process is a process for determining the differentiation-state of cells during embryonic development, angiogenesis, or development of cancer.
  - 51. A process for detecting the presence of an integrin subunit  $\alpha 10$ , or of a homologue or fragment of said integrin subunit having similar biological activity, on cells, whereby a polynucleotide or oligonucleotide chosen from the group comprising a polynucleotide or oligo-

15

25

nucleotide shown in SEQ ID No. 1 is used as a marker under hybridisation conditions wherein said polynucleotide or oligonucleotide fails to hybridise to a DNA or RNA encoding an integrin subunit  $\alpha 1$ .

- 52. A process according to claim 51, whereby said cells are chosen from the group comprising chondrocytes, smooth muscle cells, endothelial cells, osteoblasts and fibroblasts.
- 53. A process according to claim 51, whereby said fragment is a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the spliced domain.
- 54. A process according to claim 53, whereby said fragment is a peptide comprising the amino acid sequence KLGFFAHKKIPEEEKREEKLEQ.
- 55. A process according to claim 53, whereby said fragment comprises the amino acid sequence from about amino acid No. 952 to about amino acid no. 986 of SEQ ID No. 1.
- 56. A process according to claim 53, whereby said fragment comprises the amino acid sequence from about amino acid No. 140 to about amino acid No. 337 of SEQ ID No. 1.
  - 57. A process according to any one of claims 43-48, which is a process for determining the differentiation-state of cells during development, in pathological conditions, in tissue regeneration or in therapeutic and physiological reparation of cartilage.
  - 58. A process according to claim 57, wherein the pathological conditions are any pathological conditions involving the integrin subunit  $\alpha 10$ .
    - 59. A process according to claim 58, whereby said pathological conditions are rheumatoid arthritis, osteoarthrosis or cancer.
- 35 60. A process according to claim 57, whereby said cells are chosen from the group comprising chondrocytes,

15

20

smooth muscle cells, endothelial cells, osteoblasts and fibroblasts.

- 61. A process for determining the differentiation-state of cells during development, in pathological conditions, in tissue regeneration and in therapeutic and physiological reparation of cartilage, whereby a polynucleotide or oligonucleotide chosen from the nucleotide sequence shown in SEQ ID No. 1 is used as a marker under hybridisation conditions wherein said polynucleotide or oligonucleotide fails to hybridise to a DNA or RNA encoding an integrin subunit  $\alpha$ 1.
- 62. A process according to claim 61, whereby said polynucleotide or oligonucleotide is a polynucleotide or oligonucleotide coding for a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the spliced domain.
- 63. A process according to claim 62, whereby said polynucleotide or oligonucleotide is a polynucleotide or oligonucleotide coding for a peptide comprising the amino acid sequence KLGFFAHKKIPEEEKREEKLEQ.
- 64. A process according to claim 62, whereby said peptide comprises the amino acid sequence from about amino acid no. 952 to about amino acid no. 986 of SEQ ID No. 1.
- 25 65. A process according to claim 62, whereby said peptide comprises the amino acid sequence from about amino acid no. 140 to about amino acid no. 337 of SEQ ID No. 1.
  - 66. A process according to claim 61, whereby said pathological conditions are any pathological conditions involving the integrin subunit  $\alpha$ 10.
    - 67. A process according to claim 66, whereby said pathological conditions are rheumatoid arthritis, osteo-arthrosis or cancer.
- 35 68. A process according to claim 66, whereby said pathological conditions are atherosclerosis or inflammation.

25

35

- 69. A process according to any one of claims 61-68, whereby said cells are chosen from the group comprising chondrocytes, smooth muscle cells, endothelial cells, osteoblasts and fibroblasts.
- 70. A pharmaceutical composition comprising as an active ingredient a pharmaceutical agent or an antibody which is capable of using an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit  $\alpha 10$  having similar biological activity, as a target molecule.
  - 71. A pharmaceutical composition according to claim 70, for use in stimulating, inhibiting or blocking the formation of cartilage, bone or blood vessels.
- 72. A pharmaceutical composition according to claim 70, for use in preventing adhesion between tendon/ligaments and the surrounding tissue after infection, inflammation and after surgical intervention where adhesion impairs the function of the tissue.
- 73. A vaccine comprising as an active ingredient an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit  $\alpha 10$ , or DNA or RNA coding for said integrin subunit  $\alpha 10$ .
  - 74. Use of the integrin subunit  $\alpha 10$  as a marker or target in transplantation of cartilage or chondrocytes.
  - 75. A method of using binding entities having the capability of binding specifically to an integrin subunit  $\alpha 10$  comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or an integrin heterodimer comprising said subunit  $\alpha 10$  and a subunit  $\beta$ , or to homologues or fragments thereof having similar biological activity, for promoting adhesion of chondrocytes and/or osteoblasts to surfaces of implants to stimulate osseointegration.
    - 76. Use of an integrin heterodimer comprising an integrin subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$

15

25

30

35

thereof, or a homologue or fragment of said integrin or subunit all having similar biological activity, as a target for anti-adhesive drugs or molecules in tendon, ligament, skeletal muscle or other tissues where adhesion impairs the function of the tissue.

- 77. A method of stimulating, inhibiting or blocking the formation of cartilage or bone, comprising administration to a subject a suitable amount of a pharmaceutical agent or an antibody which is capable of using an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit all having similar biological activity, as a target molecule.
- 78. A method of preventing adhesion between tendon/ ligaments and the surrounding tissue after infection, inflammation and after surgical intervention where adhesion impairs the function of the tissue, comprising administration to a subject a suitable amount of a pharmaceutical agent or an antibody which is capable of using a 20 integrin heterodimer comprising a subunit  $\alpha$ 10 and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit  $\alpha 10$  having similar biological activity, as a target molecule.
  - 79. A method of stimulating extracellular matrix synthesis and repair by activation or blockage of an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or of the subunit  $\alpha 10$  thereof, or of a homologue or fragment of said integrin or subunit lpha 10 having similar biological activity.
  - 80. A method of in vitro detecting the presence of integrin binding entities, comprising interaction of an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit, with a sample, thereby causing said integrin, subunit  $\alpha 10$ , or homologue or fragment thereof having similar biological activity,

20

25

30

to modulate the binding to its natural ligand or other integrin binding proteins present in said sample.

- 81. A method of in vitro studying consequences of the interaction of a human heterodimer integrin comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit, with an integrin binding entity and thereby initiate a cellular reaction.
- 82. A method according to claim 81, whereby the con-10 sequences of said interactions are measured as alterations in cellular functions.
  - 83. A method of using DNA or RNA encoding an integrin subunit  $\alpha 10$  or homologues or fragments thereof as a target molecule.
  - 84. A method according to claim 83, whereby a polynucleotide or oligonucleotide hybridises to the DNA or RNA encoding an integrin subunit  $\alpha 10$  or homologues or fragments thereof and whereby said polynucleotide or oligonucleotide fails to hybridise to a DNA or RNA encoding en integrin subunit  $\alpha 1$ .
  - 85. A method of using a human heterodimer integrin comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit, or a DNA or RNA encoding an integrin subunit  $\alpha 10$  or homologues or fragments thereof, as a marker or target molecule during angiogenesis.
  - 86. A pharmaceutical composition comprising as an active ingredient a pharmaceutical agent or an antibody which is capable of stimulating cell surface expression of an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit  $\alpha 10$  having similar biological activity.




FIGURE 1


2/22


	Peptide	Amino acid sequence
-		
	1	DNTAQTSAYIQYEPHHSI
	2	GPGHWDR
	3	AAFDGSGQR
	4	FAMGALPD
	5	FTASLDEWTTAAR
	6	VDASFRPQGXLAP
		<u>.</u>

WO 99/51639

PCT/SE99/00544

3/22








FIGURE 3

4/22




FIGURE 4





FIGURE 5


		'	
$\begin{array}{cccccccccccccccccccccccccccccccccccc$		CATECTOCICCADGAT TOCTGCTGCCTCCATGCCACATGCCCTCAGCTACTTTGGCCGAAGTGTGGATGGT H P A Q R L A A A S M P H A L S Y P G R S V D G	1877 395
GTCTCTCCTCCCCTTTAACCTCGATGAACATCACCCCACGCCTATTCCCAGGGCCACCAGAAGCTGAAT	77 146° 19	COSCINGATORICAGORIGATECTOCOTOGOTOGOTOCCCCAGGGGGAGCCATCCTCCCAGC	1944 619
GGATALAGTGTCTTACALCATGTTGGGGGGGGGACACCATGGGTGGGGGGGGCGCCCCCGGGAGGGGGGGG	CT 216	SCEEGGCCATTGTCCATCTCACCCATCACTGGGGGGCCCCACGGCCATCAGTGGGTTCAGAGGGCAC  3 R P 1 V H L T P 5 L E V T P Q A 1 S V V Q R D	2016 643
TCAGGGAGGGGAGGGGACGTTIATCCCTCGCTGTAGGGGGGCECCAAATGCCCCATGTGCCAAGG		TGTAGGGGGGGGGCAAGAAGAGGGGGGGGGGGGGGGGGG	2088 667
CACTINGGGATACCACIGGGATATTCACCCACTGGGATATGCACCIGGGGATGTCTCTGT	TA 360 91	GREAT DE THE TENENT OF THE TEN	2106 691
CAGALACATOGECCATTTCATGCCCTGTCCCCCTCTCTCTCTCTCTCTCTCT		CATTIGATEGETETGGCEAGAGTTGTCCCCTCGGAGCCTCAGTCTGGGGAATGTCACTTGTGAG A P D G S G Q R L S P R R L R L S V G N V T C E	2232 715
AGTICIGGEATATGTGCCCGTGTGGATGCTTCATTCCAGCCTCAGGGAGCCTGGCCACCCCCACCCA	≅ \$0¢ 139	CASCTACACTICCATGGGTGGGTGGATTGACTGGACCTTGACTTGCCTTG Q L H F H V L D T 'S D Y L B P V A L T V T F A L	2304
TOCCCAMENTACATGGATGTTCTCATTGTCTTCCATGGCTCCAMENGCATCTACCCCTGGTCTGAMGTTCC C P T Y H.D V V I V L D G S H S I Y P W S Z V.Q		GACANTACTACARAGECCETGEGCTGATGAGGGCTCACCEACCTCTATACAAAACTGGTCCCTTC D H T T K P G P V L N E G S P T S I C K L V F T	2376 763
ACCTICCTACGAGACTGGTAGGGAACTGTTTAITGACCCAGAACAGTACAGGTACAGTGGACTGGTACAGTT T F L R R L V G K L F 1 D P E 0 1 0 V G L V 0 T	AT 643 187	# # TCAMAGGATTGTGGCCCTGACANTGANTGTGTCACAGGCCTGGTGCTTCAAGTGANTATGGACATCAGGGG S K D C G P D N E C V T D L V L Q V H M D I R G	2449 787
GEGGAGAGCCCTGTACAGTGGTCCCTGGGAGATTTCCGAACGAA		TCCAGAAGCCCCATTTCTGGTTCAGGTGCCGGCGGAAGTGCTGGTATCTACAACTCTGGGACAGA S R K A P F V V R G G R R K V L V S T T L Z K R	2520 811
AMECTEAGTEGGEGGGGGGGGGGAGAAACAMAGGETGECCAGGAATAATGGTGGGCTGCAACAAGGGT		AMGGAMATGCTTACAATACGAGCETGAGTATCATCTTCTCTAGAAACCTCCACCTGGCCAGTETCACTCTC K & N A Y N T S L S I I P S R N L II L A S L T P	2592 835
ASTCAGTECCATGGGGCCGACCCGAGGCTGCCAGGCTACTGGTGGGTG		CAGAGAGAGACCCAATAAAGGTGGAATGTGCCCCCCCTTCTCATCCCCCCCC	2664 859
GENERAGRACTICCTOCAGCACTARAGOCCTGTGAGGCTGGAAGAGTEALAGGCTATGGGTTGCAGTCCTGG E E L P A A L K A C E A G R V T R Y G I A V L		CCTGTCTTCCAGACTGGAGCCAAGGTGACCTTTTCCTGGAGCTTTGGGTTTAGCTGCTCCTCCTCCTCCTGAGC	2736 883
GTCACTACCTCCCCCCCCACCCACCTCCTTCCTCACACAATTMCACCTATTCCCACTCATCACC	EA 1008 307	CAGGETTTOCANCETCACTCCACCACTCACACCATCCACACAAAACCCCTTCAAGAAAACACCCCQ V F G K L T A S S D S L E R N G T L Q E N T A	2808
CATGACGATTCTTCTACATGTCACAGATGAGGCTCCTCTGACTGA	25 1090 331	CAGACCTERACCTACATOCAAFATGAGCCCCACCTCCTGTTCTCTAGTGAGTCTACCCCTALACCCCTATGAG Q T 9 A Y 1 Q Y E P H L L F 5 S E 3 T L E R Y E	2890 931
ATTITICACIONAL CAMBONICO DE LA SOLO IL POR LA CAMBONICA DE LA SOLO DEL	TC 1152	GITCACCATATOGACCCTCCCAGTGCGGCCCAGATTCAAAACCACTCTCAGGGTTCAGACCTA V H Y Y G T L P V G P G P E F K T T L R V Q N E	2952 . 955
TCCACTCATCGCTAAAGGATGGGATCTTTTTGGGATGGTGCGGGGGGGG		GOCTOCTATOGGCCATCATCTCACCTCCCACCTGGGCCCCATGGGGCAATACTTC	3024 979
TOSTTGAAGAGACCACCTTTTCCCCCCACGAATGCCACTGGAAGACGAGTTCCCCCTGCACTGC W L E G G H R L F P P R H A L B D E F P P A L Q	NG 1296 403	CTATGACTGTCTCAAGTCATCACTACAATGCAAGCTCCAGTGAACCTGACTGA	3096 1003
AACCATCCACCTACCTGGGTACTCGGTTTCTCCATGCTTTTCCGGGGGGGACCCCCCCTGTTCTCTCT	CT 1368 427	CONTROL OF THE PERSON OF THE P	3168 1027
GGGCTCCTCGATTTAGACATCGAGGAAAGTCATCCCCTTCCAGCTTAAGAAAGA		TELEMETICEOCACTGECHAGGEGETEGETETEGETATTCACCATGATTC C H L G O L A K G T E V S V G L L R L V H H E F	3240 1051
GCCCAGAGCCTCCAGGGGGGCAGAFTGGTTCATACTTTGGCAGTGAGCTCTGGCTCATTGGATACAGATA A Q S L Q G E Q I G S Y F G S E L C F L D Y D R	2G 1512 . 475	TTCCGAACACCCAAGTCCAGTCCCCGACGGGGGGGGCACCGACGACCGAACAGGGGGCACTGCC FRRAKFKSLTVVSTFELGIZEG	3312 1075
GATGEMCANCIGATGTCTTCTTCGCTCCCCCATGTTCCTGCCACCCCAGACCAACCA		CTACAGETEATTCAACCCTCCCCTTCCAGTGAGAGCTCTTCTACACCCCCCTATCCTCACCCACC	3304 1099
GTTLATGITATICTGFAGGCCAGCAGTCCTTGCTGACCCTCCAAGGACACTTCAGCCAGAACCCCCCCVVVVTLVG 0 0 5 L L T L 0 G T L 0 P E P P Q		TECCTOTOGRATECTEAN MOSE AND TECCTOGRADOUS TECCTOCCT TECCTT TECTT TECTT TECCTOCCT TECCT TE	3456 1323
GATOCTCOGTTTOCCCATGGGACCCCTTCCTGATCTGACCAAGATGGTTTTOCTCATGTGCCCCDDARFGGFAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA	75 1728 547	AACTTGCCTTCTTTCCCATALCAAATCCCTGAGGAAAAAAGAGAAGAAGAAGATGGACAATGAATG	3520 1145
GEGGGETTTTGAAEA16GCCACCACGGACCACTGTACCATGGAACCLAGAGTGGAGTAAGCC G A P L E D G H O G A L Y L Y H G T D S G V R P	CC 1800 571	0.11040.010	3600 3672
		AGGINTECTECTCOCTAGAGNICAGGETTTACCTCAGACAAGAAGAGCTGGCACCAAAACTAGCCATGCTCC CACCCTCTGCTCCTCCTGCTGATCCTGGTTCCATACCAAAACTAGGGGCCTTGTTTCGGATCTT TATTCCCAGAAGAATCAATAATTTTTTCCTCAGAAAAAAAA	3744 3816 3884

7/22



WO 99/51639

8/22



**B** IP: α10 β1

Blot: β1 β1

200 -

97 -

46 -





FIGURE 9








FIGURE 10










FIGURE 11

## Human RNA Master blot

Tissue	al0 expression	Tissue	$\alpha 10$ expression
Aorta	<del>`\+++</del>	Thyroid gland	
Trachea	÷	Salivary gland	• •
Lung	· ++	Spleen	•
Fetal lung	<del>++</del>	Fetal spleen	•
Kidney	++	Thymus	•
Fetal kidney	(+)	Fetal thymus	-
Heart	(÷)	Peripherial leucocyte	•
Fetal heart	++	Lymph node	•
Spinal cord	++	Appendix	•
Mammary gland	(+)	Placenta	•
Bone marrow	(+)	Whole brain	•
Small intestine	(+)	Fetal brain	•
Skeletal muscle	- '	Amygdala	-
Liver	. •	Caudate nucleus	-
Fetal liver	1 <b>2</b> 1	Cerebellum	-
Colon	<u>.</u>	Cerebral cortex	
Bladder	•	Frontal lobe	•
Uterus	-	Hippocampus	-
Prostate	• • • •	Medulla oblongata	-
Stomach	-	Occipitial lobe	•
Testis	-	Putamen	• .
Ovary	•	Substantia nigra	. •
Pancreas	•	Temporal lobe	-
Piutiatary gland	•	Thalamus	•
Adrenal gland	•	Subthalamic nucleus	-
			-

FIGURE 12









FIGURE 13











FIGURE 14











GGAGAGGGAGAGGGTCTGGAGTGTAGTGTATACATCACAAGATGCTCTGGGCGCCTTATCTTTATCTGCATGCCAGAAGTT 6880
ERERVWSVYTSQDALGAYLYLHARS GRGRGSGV.CIHHKMLWALIFICMPEV GEGEGLECSVYITRCSGRLSLSACQKF
CGTGGAGGAAGGCTAGGTTGCTGTCACCATACTCTCTTACTGTATTTGCATTTTATGGTGTGTGT
S W R K A R L L S P Y S L L L Y L H P M V S V G V S L R G G R L G C C H H T L S Y C I C I L W C L W V Y L S V E E G . V A V T I L S L T V P A F Y G V C G C I S
CTTGTCTGTTCTGCTCACACAGAACTCCATCTTTCCTCTTCTACTCCTGCGTCAATTCTGATACCTAGCTTCTCAA 7040
L V C S V S A H R T P S F L F Y S C V N S D T . L L N L S V L F L H T S L H L S S S T P A S I L I P S F S P C L F C F C T Q N S I F P L L L L R Q F . Y L A S Q
CCACTCACGCCCTAGTATICTTTCAAACATGACTCTAAACCTCTGGGGAGGCTACATGACCTGACTGTCTTTATTCTCC 7120
H S R P S I L F K H D S K P L G R L H D L T V F I L T T H A L V F F S N H T L N L W G G Y M T . L S L F S P L T P . Y S F Q T . L , T S G E A T . P D C L Y S P
AGTTCCTTGATCTTGTCAACCCAAGTGTTTGCTGAATGAA
Q F L D L V N P S V C . M N L . I N N A C T Y L H S S L I L S T Q V F A S . I Y K . I M L V H I Y T D D V P . S C O P K C L L N B S I N K . C L Y I F T L M
CACATTATTTTATATGTTCCGTGCCATCTAAACAGTCAAGTTGTGACTCTGTGCCAGTTTGCATGCTAGATACTGTTGGG 7280
Q I I L Y V P C H L N S Q V V T L C Q F A C . I L L G R L F Y M F R A I . T V K L . L C A S L H A R Y C W T D Y F I C S V P S K Q S S C D S V P V C M L D T V G
GAATGGTGTAGAAGACATCTGACCTCAGTGAACTGCTGACAGTGTTAATACACTATACGGGCATGCCTGCATGCA
N G V E D I . P Q . T A D S V N T L Y G H A C M Q A G M V . K T S D L S E L L T V L I H Y T G M P A C K P E W C R R H L T S V N C . Q C . Y T I R A C L H A S L
GTGTGTATGTGCATGCATATGCACACACATATGACCATATAGCATTCTTTTATCTCTCTTCTTAGCACAGAAGGGT 7440
C V Y V H A Y A H T Y I . P Y S I L L S L F L A Q K G V C M C M H M H T H T Y D H I A P F Y L S S . H R R V C V C A C I C T H I H M T I . H S F I S L L S <u>T E G</u>
TCAGTCAGTCCCGGGGGGACGACCAGAGGCCGCTAGGCTGCTGGTAGTTGTCACTGATGGAGAGTCCCATGATGGAGAG 7520
S V S P G G D D Q R P L G C W . L S L H E S P M M E R Q S V P G G T T R G R . A A G S C H . W R V P . W R P S Q S R G G R P E A A R L L V V V T D G R S H D G E
GAACTTCCAGCAGCGCTAAAGGCCTGTGAGGCTGGCAGAGTGACACGTTATGGGATTGCGGTGAGACTTGATCAAGTCCA 7600
NFQQR.RPVRLAE.HVMGLR.DLIKS GTSSSAKGL.GWQSDTLWDCGET.SSP <u>RLPXALKACEAGRVTRYGIA</u> VRLDQVQ
GTTGTTTTGTTTTGTGTGTATCCTGTGTGTGTGTGTGTGT
T L C L A T A A C A C A C A C A C A C A C A C A
GTGTGCATGCATCAGTGCACATACCATAGTGTGTATATGCGGGTCAGAGAACAACCTCAGATGTTGGTCCTCACCTTCCA 7760
V C M H Q C T Y H S V Y M R V R E Q P Q M L V L T F H C A C I S A H T I V C I C G S E N N L R C W S S P S C V H A S V H I P . C V Y A G Q R T T S D V G P H L P
TCTTGTTCCAAACTGGATATCTTGTTCACTTCGGCATACAATAAGCCAGATTAGCTGACCCACAAGTCTTGGGCAGGTCT 7840
L V P N W I S C S L R H T I S Q I S . P T S L G Q V I L F Q T G Y L V H F G I Q . A R L A D P Q V L G R S S C S K L D I L F T S A Y N K P D . L T H K S W A G L
TCTGTCTCAGCCTCCTGTCTTGGTTTGAGGCATTCTGGAATTTACAGATAAGCTTGATATCGAATTCCTGCAGCCCGG 7920
FCLSLLSLGLRHSGIYR A . YRIPAAR SVSASCLLV GILEFTDKLDIRFLQPG LSQPPVSWFEAFWNLQI5LISNSCSP
GGGATCCACTAGTTCTAGAGCGGCCGCCACCAAGGGAG 7958
GIH.F.SGRHQGS GSTSSRAAATKG GDPLVLERPPPRE

FIGURE 15f

WO 99/51639 PCT/SE99/00544




FIGURE 16




FIGURE 17

# This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

### **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:
□ BLACK BORDERS
☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
☐ FADED TEXT OR DRAWING
☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
☐ SKEWED/SLANTED IMAGES
☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
☐ GRAY SCALE DOCUMENTS
☐ LINES OR MARKS ON ORIGINAL DOCUMENT
☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

### IMAGES ARE BEST AVAILABLE COPY.

OTHER:

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.